

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C12N 15/54, 15/82, 9/10, 5/00, C12P 17/06</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/63391</b> <b>(43) International Publication Date:</b> 26 October 2000 (26.10.00)
<b>(21) International Application Number:</b> PCT/US00/10368 <b>(22) International Filing Date:</b> 14 April 2000 (14.04.00)  <b>(30) Priority Data:</b> 60/129,899                      15 April 1999 (15.04.99)                      US 60/146,461                      30 July 1999 (30.07.99)                      US  <b>(71) Applicant (for all designated States except US):</b> CALGENE LLC [US/US]; 1920 Fifth Street, Davis, CA 95616 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> SAVIDGE, Beth [US/US]; 1920 Fifth Street, Davis, CA 95616 (US). LASSNER, Michael, W. [US/US]; 1920 Fifth Street, Davis, CA 95616 (US). WEISS, James, D. [US/US]; 800 N. Lindbergh Blvd., St. Louis, MO 63167 (US). POST-BEITTMILLER, Dusty [US/US]; 800 N. Lindbergh Blvd., St. Louis, MO 63167 (US).  <b>(74) Agents:</b> SCHWEDLER, Carl, J. et al.; Calgene LLC, 1920 Fifth Street, Davis, CA 95616 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> NUCLEIC ACID SEQUENCES TO PROTEINS INVOLVED IN TOCOPHEROL SYNTHESIS  <b>(57) Abstract</b>  Nucleic acid sequences and methods are provided for producing plants and seeds having altered tocopherol content and compositions. The methods find particular use in increasing the tocopherol levels in plants, and in providing desirable tocopherol compositions in a host plant cell.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## NUCLEIC ACID SEQUENCES TO PROTEINS INVOLVED IN TOCOPHEROL SYNTHESIS

5

### INTRODUCTION

This application claims the benefit of the filing date of the provisional Application U.S. Serial Number 60/129,899, filed April 15, 1999, and the provisional Application, U.S.  
10 Serial Number 60/146,461, filed July 30, 1999.

### TECHNICAL FIELD

The present invention is directed to nucleic acid and amino acid sequences and constructs, and methods related thereto.

15

### BACKGROUND

Isoprenoids are ubiquitous compounds found in all living organisms. Plants synthesize a diverse array of greater than 22,000 isoprenoids (Connolly and Hill (1992) *Dictionary of Terpenoids*, Chapman and Hall, New York, NY). In plants, isoprenoids play essential roles in  
20 particular cell functions such as production of sterols, contributing to eukaryotic membrane architecture, acyclic polyprenoids found in the side chain of ubiquinone and plastoquinone, growth regulators like abscisic acid, gibberellins, brassinosteroids or the photosynthetic pigments chlorophylls and carotenoids. Although the physiological role of other plant isoprenoids is less evident, like that of the vast array of secondary metabolites, some are  
25 known to play key roles mediating the adaptative responses to different environmental challenges. In spite of the remarkable diversity of structure and function, all isoprenoids originate from a single metabolic precursor, isopentenyl diphosphate (IPP) (Wright, (1961) *Annu. Rev. Biochem.* 20:525-548; and Spurgeon and Porter, (1981) in Biosynthesis of Isoprenoid Compounds., Porter and Spurgeon eds (John Wiley, New York) Vol. 1, pp1-46).

30

A number of unique and interconnected biochemical pathways derived from the isoprenoid pathway leading to secondary metabolites, including tocopherols, exist in chloroplasts of higher plants. Tocopherols not only perform vital functions in plants, but are

also important from mammalian nutritional perspectives. In plastids, tocopherols account for up to 40% of the total quinone pool.

5 Tocopherols and tocotrienols (unsaturated tocopherol derivatives) are well known antioxidants, and play an important role in protecting cells from free radical damage, and in the prevention of many diseases, including cardiac disease, cancer, cataracts, retinopathy, Alzheimer's disease, and neurodegeneration, and have been shown to have beneficial effects on symptoms of arthritis, and in anti-aging. Vitamin E is used in chicken feed for improving the shelf life, appearance, flavor, and oxidative stability of meat, and to transfer tocopherols from feed to eggs. Vitamin E has been shown to be essential for normal reproduction, improves overall performance, and enhances immunocompetence in livestock animals. Vitamin E  
10 supplement in animal feed also imparts oxidative stability to milk products.

The demand for natural tocopherols as supplements has been steadily growing at a rate of 10-20% for the past three years. At present, the demand exceeds the supply for natural tocopherols, which are known to be more biopotent than racemic mixtures of synthetically  
15 produced tocopherols. Naturally occurring tocopherols are all *d*-stereoisomers, whereas synthetic  $\alpha$ -tocopherol is a mixture of eight *d,l*- $\alpha$ -tocopherol isomers, only one of which (12.5%) is identical to the natural *d*- $\alpha$ -tocopherol. Natural *d*- $\alpha$ -tocopherol has the highest vitamin E activity (1.49 IU/mg) when compared to other natural tocopherols or tocotrienols. The synthetic  $\alpha$ -tocopherol has a vitamin E activity of 1.1 IU/mg. In 1995, the worldwide  
20 market for raw refined tocopherols was \$1020 million; synthetic materials comprised 85-88% of the market, the remaining 12-15% being natural materials. The best sources of natural tocopherols and tocotrienols are vegetable oils and grain products. Currently, most of the natural Vitamin E is produced from  $\gamma$ -tocopherol derived from soy oil processing, which is subsequently converted to  $\alpha$ -tocopherol by chemical modification ( $\alpha$ -tocopherol exhibits the  
25 greatest biological activity).

Methods of enhancing the levels of tocopherols and tocotrienols in plants, especially levels of the more desirable compounds that can be used directly, without chemical modification, would be useful to the art as such molecules exhibit better functionality and bioavailability.

30 In addition, methods for the increased production of other isoprenoid derived compounds in a host plant cell is desirable. Furthermore, methods for the production of particular isoprenoid compounds in a host plant cell is also needed.

## SUMMARY OF THE INVENTION

5           The present invention is directed to prenyltransferase (PT), and in particular to PT polynucleotides and polypeptides. The polynucleotides and polypeptides of the present invention include those derived from prokaryotic and eukaryotic sources.

          Thus, one aspect of the present invention relates to isolated polynucleotide sequences encoding prenyltransferase proteins. In particular, isolated nucleic acid sequences encoding  
10   PT proteins from bacterial and plant sources are provided.

          Another aspect of the present invention relates to oligonucleotides which include partial or complete PT encoding sequences.

          It is also an aspect of the present invention to provide recombinant DNA constructs which can be used for transcription or transcription and translation (expression) of  
15   prenyltransferase. In particular, constructs are provided which are capable of transcription or transcription and translation in host cells.

          In another aspect of the present invention, methods are provided for production of prenyltransferase in a host cell or progeny thereof. In particular, host cells are transformed or transfected with a DNA construct which can be used for transcription or transcription and  
20   translation of prenyltransferase. The recombinant cells which contain prenyltransferase are also part of the present invention.

          In a further aspect, the present invention relates to methods of using polynucleotide and polypeptide sequences to modify the tocopherol content of host cells, particularly in host plant cells. Plant cells having such a modified tocopherol content are also contemplated  
25   herein.

          The modified plants, seeds and oils obtained by the expression of the prenyltransferases are also considered part of the invention.

## BRIEF DESCRIPTION OF THE DRAWINGS

30           Figure 1 provides an amino acid sequence alignment between ATPT2, ATPT3, ATPT4, ATPT8, and ATPT12 are performed using ClustalW.

Figure 2 provides a schematic picture of the expression construct pCGN10800.

Figure 3 provides a schematic picture of the expression construct pCGN10801.

Figure 4 provides a schematic picture of the expression construct pCGN10803.

Figure 5 provides a schematic picture of the expression construct pCGN10806.

5 Figure 6 provides a schematic picture of the expression construct pCGN10807.

Figure 7 provides a schematic picture of the expression construct pCGN10808.

Figure 8 provides a schematic picture of the expression construct pCGN10809.

Figure 9 provides a schematic picture of the expression construct pCGN10810.

Figure 10 provides a schematic picture of the expression construct pCGN10811.

10 Figure 11 provides a schematic picture of the expression construct pCGN10812.

Figure 12 provides a schematic picture of the expression construct pCGN10813.

Figure 13 provides a schematic picture of the expression construct pCGN10814.

Figure 14 provides a schematic picture of the expression construct pCGN10815.

Figure 15 provides a schematic picture of the expression construct pCGN10816.

15 Figure 16 provides a schematic picture of the expression construct pCGN10817.

Figure 17 provides a schematic picture of the expression construct pCGN10819.

Figure 18 provides a schematic picture of the expression construct pCGN10824.

Figure 19 provides a schematic picture of the expression construct pCGN10825.

Figure 20 provides a schematic picture of the expression construct pCGN10826.

20 Figure 21 provides an amino acid sequence alignment using ClustalW between the *Synechocystis* sequence knockouts.

Figure 22 provides an amino acid sequence of the ATPT2, ATPT3, ATPT4, ATPT8, and ATPT12 protein sequences from *Arabidopsis* and the slr1736, slr0926, slr1899, slr0056, and the slr1518 amino acid sequences from *Synechocystis*.

25 Figure 23 provides the results of the enzymatic assay from preparations of wild type *Synechocystis* strain 6803, and *Synechocystis* slr1736 knockout.

Figure 24 provides bar graphs of HPLC data obtained from seed extracts of transgenic *Arabidopsis* containing pCGN10822, which provides of the expression of the ATPT2 sequence, in the sense orientation, from the napin promoter. Provided are graphs for alpha, gamma, and delta tocopherols, as well as total tocopherol for 22 transformed lines, as well as a nontransformed (wildtype) control.

30

Figure 25 provides a bar graph of HPLC analysis of seed extracts from *Arabidopsis* plants transformed with pCGN10803 (35S-ATPT2, in the antisense orientation), pCGN10802

(line 1625, napin ATPT2 in the sense orientation), pCGN10809 (line 1627, 35S-ATPT3 in the sense orientation), a nontransformed (wt) control, and a empty vector transformed control.

5

## DETAILED DESCRIPTION OF THE INVENTION

The present invention provides, *inter alia*, compositions and methods for altering (for example, increasing and decreasing) the tocopherol levels and/or modulating their ratios in  
10 host cells. In particular, the present invention provides polynucleotides, polypeptides, and methods of use thereof for the modulation of tocopherol content in host plant cells.

The present invention provides polynucleotide and polypeptide sequences involved in the prenylation of straight chain and aromatic compounds. Straight chain prenyl transferases as used herein comprises sequences which encode proteins involved in the prenylation of  
15 straight chain compounds, including, but not limited to, geranyl geranyl pyrophosphate and farnesyl pyrophosphate. Aromatic prenyl transferases, as used herein, comprises sequences which encode proteins involved in the prenylation of aromatic compounds, including, but not limited to, menaquinone, ubiquinone, chlorophyll, and homogentisic acid. The prenyl transferase of the present invention preferably prenylates homogentisic acid.

20 The biosynthesis of  $\alpha$ -tocopherol in higher plants involves condensation of homogentisic acid and phytylpyrophosphate to form 2-methyl-6 phytylbenzoquinol that can, by cyclization and subsequent methylations (Fiedler et al., 1982, *Planta*, 155: 511-515, Soll et al., 1980, *Arch. Biochem. Biophys.* 204: 544-550, Marshall et al., 1985 *Phytochem.*, 24: 1705-1711, all of which are herein incorporated by reference in their entirety), form various  
25 tocopherols. The *Arabidopsis pds2* mutant identified and characterized by Norris *et al.* (1995), is deficient in tocopherol and plastoquinone-9 accumulation. Further genetic and biochemical analysis suggests that the protein encoded by *PDS2* may be responsible for the prenylation of homogentisic acid. This may be a rate limiting step in tocopherol biosynthesis, and this gene has yet to be isolated. Thus, it is an aspect of the present invention to provide  
30 polynucleotides and polypeptides involved in the prenylation of homogentisic acid.

### Isolated Polynucleotides, Proteins, and Polypeptides



A first aspect of the present invention relates to isolated prenyltransferase polynucleotides. The polynucleotide sequences of the present invention include isolated polynucleotides that encode the polypeptides of the invention having a deduced amino acid sequence selected from the group of sequences set forth in the Sequence Listing and to other polynucleotide sequences closely related to such sequences and variants thereof.

The invention provides a polynucleotide sequence identical over its entire length to each coding sequence as set forth in the Sequence Listing. The invention also provides the coding sequence for the mature polypeptide or a fragment thereof, as well as the coding sequence for the mature polypeptide or a fragment thereof in a reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, pro-, or prepro- protein sequence. The polynucleotide can also include non-coding sequences, including for example, but not limited to, non-coding 5' and 3' sequences, such as the transcribed, untranslated sequences, termination signals, ribosome binding sites, sequences that stabilize mRNA, introns, polyadenylation signals, and additional coding sequence that encodes additional amino acids. For example, a marker sequence can be included to facilitate the purification of the fused polypeptide. Polynucleotides of the present invention also include polynucleotides comprising a structural gene and the naturally associated sequences that control gene expression.

The invention also includes polynucleotides of the formula:



wherein, at the 5' end, X is hydrogen, and at the 3' end, Y is hydrogen or a metal,  $R_1$  and  $R_3$  are any nucleic acid residue,  $n$  is an integer between 1 and 3000, preferably between 1 and 1000 and  $R_2$  is a nucleic acid sequence of the invention, particularly a nucleic acid sequence selected from the group set forth in the Sequence Listing and preferably those of SEQ ID NOs: 1, 3, 5, 7, 8, 10, 11, 13-16, 18, 23, 29, 36, and 38. In the formula,  $R_2$  is oriented so that its 5' end residue is at the left, bound to  $R_1$ , and its 3' end residue is at the right, bound to  $R_3$ . Any stretch of nucleic acid residues denoted by either R group, where R is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer.

The invention also relates to variants of the polynucleotides described herein that encode for variants of the polypeptides of the invention. Variants that are fragments of the polynucleotides of the invention can be used to synthesize full-length polynucleotides of the invention. Preferred embodiments are polynucleotides encoding polypeptide variants wherein

5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues of a polypeptide sequence of the invention are substituted, added or deleted, in any combination. Particularly preferred are substitutions, additions, and deletions that are silent such that they do not alter the properties or activities of the polynucleotide or polypeptide.

5 Further preferred embodiments of the invention that are at least 50%, 60%, or 70% identical over their entire length to a polynucleotide encoding a polypeptide of the invention, and polynucleotides that are complementary to such polynucleotides. More preferable are polynucleotides that comprise a region that is at least 80% identical over its entire length to a polynucleotide encoding a polypeptide of the invention and polynucleotides that are  
10 complementary thereto. In this regard, polynucleotides at least 90% identical over their entire length are particularly preferred, those at least 95% identical are especially preferred. Further, those with at least 97% identity are highly preferred and those with at least 98% and 99% identity are particularly highly preferred, with those at least 99% being the most highly preferred.

15 Preferred embodiments are polynucleotides that encode polypeptides that retain substantially the same biological function or activity as the mature polypeptides encoded by the polynucleotides set forth in the Sequence Listing.

The invention further relates to polynucleotides that hybridize to the above-described sequences. In particular, the invention relates to polynucleotides that hybridize under  
20 stringent conditions to the above-described polynucleotides. As used herein, the terms "stringent conditions" and "stringent hybridization conditions" mean that hybridization will generally occur if there is at least 95% and preferably at least 97% identity between the sequences. An example of stringent hybridization conditions is overnight incubation at 42°C in a solution comprising 50% formamide, 5x SSC (150 mM NaCl, 15 mM trisodium citrate),  
25 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/milliliter denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1x SSC at approximately 65°C. Other hybridization and wash conditions are well known and are exemplified in Sambrook, *et al.*, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, NY (1989), particularly Chapter 11.

30 The invention also provides a polynucleotide consisting essentially of a polynucleotide sequence obtainable by screening an appropriate library containing the complete gene for a polynucleotide sequence set forth in the Sequence Listing under stringent hybridization conditions with a probe having the sequence of said polynucleotide sequence or

a fragment thereof; and isolating said polynucleotide sequence. Fragments useful for obtaining such a polynucleotide include, for example, probes and primers as described herein.

As discussed herein regarding polynucleotide assays of the invention, for example, polynucleotides of the invention can be used as a hybridization probe for RNA, cDNA, or genomic DNA to isolate full length cDNAs or genomic clones encoding a polypeptide and to isolate cDNA or genomic clones of other genes that have a high sequence similarity to a polynucleotide set forth in the Sequence Listing. Such probes will generally comprise at least 15 bases. Preferably such probes will have at least 30 bases and can have at least 50 bases. Particularly preferred probes will have between 30 bases and 50 bases, inclusive.

The coding region of each gene that comprises or is comprised by a polynucleotide sequence set forth in the Sequence Listing may be isolated by screening using a DNA sequence provided in the Sequence Listing to synthesize an oligonucleotide probe. A labeled oligonucleotide having a sequence complementary to that of a gene of the invention is then used to screen a library of cDNA, genomic DNA or mRNA to identify members of the library which hybridize to the probe. For example, synthetic oligonucleotides are prepared which correspond to the prenyltransferase EST sequences. The oligonucleotides are used as primers in polymerase chain reaction (PCR) techniques to obtain 5' and 3' terminal sequence of prenyl transferase genes. Alternatively, where oligonucleotides of low degeneracy can be prepared from particular prenyltransferase peptides, such probes may be used directly to screen gene libraries for prenyltransferase gene sequences. In particular, screening of cDNA libraries in phage vectors is useful in such methods due to lower levels of background hybridization.

Typically, a prenyltransferase sequence obtainable from the use of nucleic acid probes will show 60-70% sequence identity between the target prenyltransferase sequence and the encoding sequence used as a probe. However, lengthy sequences with as little as 50-60% sequence identity may also be obtained. The nucleic acid probes may be a lengthy fragment of the nucleic acid sequence, or may also be a shorter, oligonucleotide probe. When longer nucleic acid fragments are employed as probes (greater than about 100 bp), one may screen at lower stringencies in order to obtain sequences from the target sample which have 20-50% deviation (i.e., 50-80% sequence homology) from the sequences used as probe.

Oligonucleotide probes can be considerably shorter than the entire nucleic acid sequence encoding an prenyltransferase enzyme, but should be at least about 10, preferably at least about 15, and more preferably at least about 20 nucleotides. A higher degree of sequence

identity is desired when shorter regions are used as opposed to longer regions. It may thus be desirable to identify regions of highly conserved amino acid sequence to design oligonucleotide probes for detecting and recovering other related prenyltransferase genes. Shorter probes are often particularly useful for polymerase chain reactions (PCR), especially when highly conserved sequences can be identified. (See, Gould, *et al.*, *PNAS USA* (1989) 86:1934-1938.).

Another aspect of the present invention relates to prenyltransferase polypeptides. Such polypeptides include isolated polypeptides set forth in the Sequence Listing, as well as polypeptides and fragments thereof, particularly those polypeptides which exhibit prenyltransferase activity and also those polypeptides which have at least 50%, 60% or 70% identity, preferably at least 80% identity, more preferably at least 90% identity, and most preferably at least 95% identity to a polypeptide sequence selected from the group of sequences set forth in the Sequence Listing, and also include portions of such polypeptides, wherein such portion of the polypeptide preferably includes at least 30 amino acids and more preferably includes at least 50 amino acids.

"Identity", as is well understood in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as determined by the match between strings of such sequences. "Identity" can be readily calculated by known methods including, but not limited to, those described in *Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York (1988); *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data, Part I*, Griffin, A.M. and Griffin, H.G., eds., Humana Press, New Jersey (1994); *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press (1987); *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., Stockton Press, New York (1991); and Carillo, H., and Lipman, D., *SIAM J Applied Math*, 48:1073 (1988). Methods to determine identity are designed to give the largest match between the sequences tested. Moreover, methods to determine identity are codified in publicly available programs. Computer programs which can be used to determine identity between two sequences include, but are not limited to, GCG (Devereux, J., et al., *Nucleic Acids Research* 12(1):387 (1984); suite of five BLAST programs, three designed for nucleotide sequences queries (BLASTN, BLASTX, and TBLASTX) and two designed for protein sequence queries (BLASTP and TBLASTN)

(Coulson, *Trends in Biotechnology*, 12: 76-80 (1994); Birren, *et al.*, *Genome Analysis*, 1: 543-559 (1997)). The BLAST X program is publicly available from NCBI and other sources (*BLAST Manual*, Altschul, S., *et al.*, NCBI NLM NIH, Bethesda, MD 20894; Altschul, S., *et al.*, *J. Mol. Biol.*, 215:403-410 (1990)). The well known Smith Waterman algorithm can also  
 5 be used to determine identity.

Parameters for polypeptide sequence comparison typically include the following:

Algorithm: Needleman and Wunsch, *J. Mol. Biol.* 48:443-453 (1970)

Comparison matrix: BLOSSUM62 from Hentikoff and Hentikoff, *Proc. Natl. Acad. Sci USA* 89:10915-10919 (1992)

10 Gap Penalty: 12

Gap Length Penalty: 4

A program which can be used with these parameters is publicly available as the "gap" program from Genetics Computer Group, Madison Wisconsin. The above parameters along with no penalty for end gap are the default parameters for peptide comparisons.

15 Parameters for polynucleotide sequence comparison include the following:

Algorithm: Needleman and Wunsch, *J. Mol. Biol.* 48:443-453 (1970)

Comparison matrix: matches = +10; mismatches = 0

Gap Penalty: 50

Gap Length Penalty: 3

20 A program which can be used with these parameters is publicly available as the "gap" program from Genetics Computer Group, Madison Wisconsin. The above parameters are the default parameters for nucleic acid comparisons.

The invention also includes polypeptides of the formula:



25 wherein, at the amino terminus, X is hydrogen, and at the carboxyl terminus, Y is hydrogen or a metal, R<sub>1</sub> and R<sub>3</sub> are any amino acid residue, n is an integer between 1 and 1000, and R<sub>2</sub> is an amino acid sequence of the invention, particularly an amino acid sequence selected from the group set forth in the Sequence Listing and preferably those encoded by the sequences provided in SEQ ID NOs: 2, 4, 6, 9, 12, 17, 19-22, 24-28, 30, 32-35, 37, and 39. In the  
 30 formula, R<sub>2</sub> is oriented so that its amino terminal residue is at the left, bound to R<sub>1</sub>, and its carboxy terminal residue is at the right, bound to R<sub>3</sub>. Any stretch of amino acid residues denoted by either R group, where R is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer.

Polypeptides of the present invention include isolated polypeptides encoded by a polynucleotide comprising a sequence selected from the group of a sequence contained in the Sequence Listing set forth herein .

5 The polypeptides of the present invention can be mature protein or can be part of a fusion protein.

Fragments and variants of the polypeptides are also considered to be a part of the invention. A fragment is a variant polypeptide which has an amino acid sequence that is entirely the same as part but not all of the amino acid sequence of the previously described polypeptides. The fragments can be "free-standing" or comprised within a larger polypeptide  
10 of which the fragment forms a part or a region, most preferably as a single continuous region. Preferred fragments are biologically active fragments which are those fragments that mediate activities of the polypeptides of the invention, including those with similar activity or improved activity or with a decreased activity. Also included are those fragments that antigenic or immunogenic in an animal, particularly a human.

15 Variants of the polypeptide also include polypeptides that vary from the sequences set forth in the Sequence Listing by conservative amino acid substitutions, substitution of a residue by another with like characteristics. In general, such substitutions are among Ala, Val, Leu and Ile; between Ser and Thr; between Asp and Glu; between Asn and Gln; between Lys and Arg; or between Phe and Tyr. Particularly preferred are variants in which 5 to 10; 1  
20 to 5; 1 to 3 or one amino acid(s) are substituted, deleted, or added, in any combination.

Variants that are fragments of the polypeptides of the invention can be used to produce the corresponding full length polypeptide by peptide synthesis. Therefore, these variants can be used as intermediates for producing the full-length polypeptides of the invention.

25 The polynucleotides and polypeptides of the invention can be used, for example, in the transformation of host cells, such as plant host cells, as further discussed herein.

The invention also provides polynucleotides that encode a polypeptide that is a mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids within the mature polypeptide (for example, when the mature form of the protein has more than one  
30 polypeptide chain). Such sequences can, for example, play a role in the processing of a protein from a precursor to a mature form, allow protein transport, shorten or lengthen protein half-life, or facilitate manipulation of the protein in assays or production. It is contemplated



that cellular enzymes can be used to remove any additional amino acids from the mature protein.

A precursor protein, having the mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. The inactive precursors generally are activated when the prosequences are removed. Some or all of the prosequences may be removed prior to activation. Such precursor protein are generally called proproteins.

### **Plant Constructs and Methods of Use**

Of particular interest is the use of the nucleotide sequences in recombinant DNA constructs to direct the transcription or transcription and translation (expression) of the prenyltransferase sequences of the present invention in a host plant cell. The expression constructs generally comprise a promoter functional in a host plant cell operably linked to a nucleic acid sequence encoding a prenyltransferase of the present invention and a transcriptional termination region functional in a host plant cell.

A first nucleic acid sequence is "operably linked" or "operably associated" with a second nucleic acid sequence when the sequences are so arranged that the first nucleic acid sequence affects the function of the second nucleic-acid sequence. Preferably, the two sequences are part of a single contiguous nucleic acid molecule and more preferably are adjacent. For example, a promoter is operably linked to a gene if the promoter regulates or mediates transcription of the gene in a cell.

Those skilled in the art will recognize that there are a number of promoters which are functional in plant cells, and have been described in the literature. Chloroplast and plastid specific promoters, chloroplast or plastid functional promoters, and chloroplast or plastid operable promoters are also envisioned.

One set of plant functional promoters are constitutive promoters such as the CaMV35S or FMV35S promoters that yield high levels of expression in most plant organs. Enhanced or duplicated versions of the CaMV35S and FMV35S promoters are useful in the practice of this invention (Odell, *et al.* (1985) *Nature* 313:810-812; Rogers, U.S. Patent Number 5,378, 619). In addition, it may also be preferred to bring about expression of the prenyltransferase gene in specific tissues of the plant, such as leaf, stem, root, tuber, seed, fruit, etc., and the promoter chosen should have the desired tissue and developmental specificity.

Of particular interest is the expression of the nucleic acid sequences of the present invention from transcription initiation regions which are preferentially expressed in a plant seed tissue. Examples of such seed preferential transcription initiation sequences include those sequences derived from sequences encoding plant storage protein genes or from genes involved in fatty acid biosynthesis in oilseeds. Examples of such promoters include the 5' regulatory regions from such genes as napin (Kridl *et al.*, *Seed Sci. Res.* 1:209:219 (1991)), phaseolin, zein, soybean trypsin inhibitor, ACP, stearyl-ACP desaturase, soybean  $\alpha'$  subunit of  $\beta$ -conglycinin (soy 7s, (Chen *et al.*, *Proc. Natl. Acad. Sci.*, 83:8560-8564 (1986))) and oleosin.

It may be advantageous to direct the localization of proteins conferring prenyltransferase to a particular subcellular compartment, for example, to the mitochondrion, endoplasmic reticulum, vacuoles, chloroplast or other plastidic compartment. For example, where the genes of interest of the present invention will be targeted to plastids, such as chloroplasts, for expression, the constructs will also employ the use of sequences to direct the gene to the plastid. Such sequences are referred to herein as chloroplast transit peptides (CTP) or plastid transit peptides (PTP). In this manner, where the gene of interest is not directly inserted into the plastid, the expression construct will additionally contain a gene encoding a transit peptide to direct the gene of interest to the plastid. The chloroplast transit peptides may be derived from the gene of interest, or may be derived from a heterologous sequence having a CTP. Such transit peptides are known in the art. See, for example, Von Heijne *et al.* (1991) *Plant Mol. Biol. Rep.* 9:104-126; Clark *et al.* (1989) *J. Biol. Chem.* 264:17544-17550; della-Cioppa *et al.* (1987) *Plant Physiol.* 84:965-968; Romer *et al.* (1993) *Biochem. Biophys. Res Commun.* 196:1414-1421; and, Shah *et al.* (1986) *Science* 233:478-481.

Depending upon the intended use, the constructs may contain the nucleic acid sequence which encodes the entire prenyltransferase protein, or a portion thereof. For example, where antisense inhibition of a given prenyltransferase protein is desired, the entire prenyltransferase sequence is not required. Furthermore, where prenyltransferase sequences used in constructs are intended for use as probes, it may be advantageous to prepare constructs containing only a particular portion of a prenyltransferase encoding sequence, for example a sequence which is discovered to encode a highly conserved prenyltransferase region.



The skilled artisan will recognize that there are various methods for the inhibition of expression of endogenous sequences in a host cell. Such methods include, but are not limited to, antisense suppression (Smith, *et al.* (1988) *Nature* 334:724-726), co-suppression (Napoli, *et al.* (1989) *Plant Cell* 2:279-289), ribozymes (PCT Publication WO 97/10328), and combinations of sense and antisense Waterhouse, *et al.* (1998) *Proc. Natl. Acad. Sci. USA* 95:13959-13964. Methods for the suppression of endogenous sequences in a host cell typically employ the transcription or transcription and translation of at least a portion of the sequence to be suppressed. Such sequences may be homologous to coding as well as non-coding regions of the endogenous sequence.

Regulatory transcript termination regions may be provided in plant expression constructs of this invention as well. Transcript termination regions may be provided by the DNA sequence encoding the prenyltransferase or a convenient transcription termination region derived from a different gene source, for example, the transcript termination region which is naturally associated with the transcript initiation region. The skilled artisan will recognize that any convenient transcript termination region which is capable of terminating transcription in a plant cell may be employed in the constructs of the present invention.

Alternatively, constructs may be prepared to direct the expression of the prenyltransferase sequences directly from the host plant cell plastid. Such constructs and methods are known in the art and are generally described, for example, in Svab, *et al.* (1990) *Proc. Natl. Acad. Sci. USA* 87:8526-8530 and Svab and Maliga (1993) *Proc. Natl. Acad. Sci. USA* 90:913-917 and in U.S. Patent Number 5,693,507.

The prenyltransferase constructs of the present invention can be used in transformation methods with additional constructs providing for the expression of other nucleic acid sequences encoding proteins involved in the production of tocopherols, or tocopherol precursors such as homogentisic acid and/or phytylpyrophosphate. Nucleic acid sequences encoding proteins involved in the production of homogentisic acid are known in the art, and include but not are limited to, 4-hydroxyphenylpyruvate dioxygenase (HPPD, EC 1.13.11.27) described for example, by Garcia, *et al.* ((1999) *Plant Physiol.* 119(4):1507-1516), mono or bifunctional tyrA (described for example by Xia, *et al.* (1992) *J. Gen Microbiol.* 138:1309-1316, and Hudson, *et al.* (1984) *J. Mol. Biol.* 180:1023-1051), Oxygenase, 4-hydroxyphenylpyruvate di- (9CI), 4-Hydroxyphenylpyruvate dioxygenase; p-Hydroxyphenylpyruvate dioxygenase; p-Hydroxyphenylpyruvate hydroxylase; p-Hydroxyphenylpyruvate oxidase; p-Hydroxyphenylpyruvic acid hydroxylase; p-

Hydroxyphenylpyruvic hydroxylase; p-Hydroxyphenylpyruvic oxidase), 4-hydroxyphenylacetate, NAD(P)H: oxygen oxidoreductase (1-hydroxylating); 4-hydroxyphenylacetate 1-monooxygenase, and the like. In addition, constructs for the expression of nucleic acid sequences encoding proteins involved in the production of phytylpyrophosphate can also be employed with the prenyltransferase constructs of the present invention. Nucleic acid sequences encoding proteins involved in the production of phytylpyrophosphate are known in the art, and include, but are not limited to geranylgeranylpyrophosphate synthase (GGPPS), geranylgeranylpyrophosphate reductase (GGH), 1-deoxyxylulose-5-phosphate synthase, 1-deoxy-D-xylulose-5-phosphate reductoisomerase, 4-diphosphocytidyl-2-C-methylerythritol synthase, isopentyl pyrophosphate isomerase.

The prenyltransferase sequences of the present invention find use in the preparation of transformation constructs having a second expression cassette for the expression of additional sequences involved in tocopherol biosynthesis. Additional tocopherol biosynthesis sequences of interest in the present invention include, but are not limited to gamma-tocopherol methyltransferase (Shintani, *et al.* (1998) *Science* 282(5396):2098-2100), tocopherol cyclase, and tocopherol methyltransferase.

A plant cell, tissue, organ, or plant into which the recombinant DNA constructs containing the expression constructs have been introduced is considered transformed, transfected, or transgenic. A transgenic or transformed cell or plant also includes progeny of the cell or plant and progeny produced from a breeding program employing such a transgenic plant as a parent in a cross and exhibiting an altered phenotype resulting from the presence of a prenyltransferase nucleic acid sequence.

Plant expression or transcription constructs having a prenyltransferase as the DNA sequence of interest for increased or decreased expression thereof may be employed with a wide variety of plant life, particularly, plant life involved in the production of vegetable oils for edible and industrial uses. Particularly preferred plants for use in the methods of the present invention include, but are not limited to: *Acacia*, alfalfa, aneth, apple, apricot, artichoke, arugula, asparagus, avocado, banana, barley, beans, beet, blackberry, blueberry, broccoli, brussels sprouts, cabbage, canola, cantaloupe, carrot, cassava, cauliflower, celery, cherry, chicory, cilantro, citrus, clementines, coffee, corn, cotton, cucumber, Douglas fir, eggplant, endive, escarole, eucalyptus, fennel, figs, garlic, gourd, grape, grapefruit, honey dew, jicama, kiwifruit, lettuce, leeks, lemon, lime, Loblolly pine, mango, melon, mushroom,

nectarine, nut, oat, oil palm, oil seed rape, okra, onion, orange, an ornamental plant, papaya, parsley, pea, peach, peanut, pear, pepper, persimmon, pine, pineapple, plantain, plum, pomegranate, poplar, potato, pumpkin, quince, radiata pine, radicchio, radish, raspberry, rice, rye, sorghum, Southern pine, soybean, spinach, squash, strawberry, sugarbeet, sugarcane, sunflower, sweet potato, sweetgum, tangerine, tea, tobacco, tomato, tritcale, turf, turnip, a vine, watermelon, wheat, yams, and zucchini.

Most especially preferred are temperate oilseed crops. Temperate oilseed crops of interest include, but are not limited to, rapeseed (Canola and High Erucic Acid varieties), sunflower, safflower, cotton, soybean, peanut, coconut and oil palms, and corn. Depending on the method for introducing the recombinant constructs into the host cell, other DNA sequences may be required. Importantly, this invention is applicable to dicotyledons and monocotyledons species alike and will be readily applicable to new and/or improved transformation and regulation techniques.

Of particular interest, is the use of prenyltransferase constructs in plants to produce plants or plant parts, including, but not limited to leaves, stems, roots, reproductive, and seed, with a modified content of tocopherols in plant parts having transformed plant cells.

For immunological screening, antibodies to the protein can be prepared by injecting rabbits or mice with the purified protein or portion thereof, such methods of preparing antibodies being well known to those in the art. Either monoclonal or polyclonal antibodies can be produced, although typically polyclonal antibodies are more useful for gene isolation. Western analysis may be conducted to determine that a related protein is present in a crude extract of the desired plant species, as determined by cross-reaction with the antibodies to the encoded proteins. When cross-reactivity is observed, genes encoding the related proteins are isolated by screening expression libraries representing the desired plant species. Expression libraries can be constructed in a variety of commercially available vectors, including lambda gt11, as described in Sambrook, *et al.* (*Molecular Cloning: A Laboratory Manual*, Second Edition (1989) Cold Spring Harbor Laboratory, Cold Spring Harbor, New York).

To confirm the activity and specificity of the proteins encoded by the identified nucleic acid sequences as prenyltransferase enzymes, *in vitro* assays are performed in insect cell cultures using baculovirus expression systems. Such baculovirus expression systems are known in the art and are described by Lee, *et al.* U.S. Patent Number 5,348,886, the entirety of which is herein incorporated by reference.

In addition, other expression constructs may be prepared to assay for protein activity utilizing different expression systems. Such expression constructs are transformed into yeast or prokaryotic host and assayed for prenyltransferase activity. Such expression systems are known in the art and are readily available through commercial sources.

5 In addition to the sequences described in the present invention, DNA coding sequences useful in the present invention can be derived from algae, fungi, bacteria, mammalian sources, plants, etc. Homology searches in existing databases using signature sequences corresponding to conserved nucleotide and amino acid sequences of prenyltransferase can be employed to isolate equivalent, related genes from other sources such as plants and  
10 microorganisms. Searches in EST databases can also be employed. Furthermore, the use of DNA sequences encoding enzymes functionally enzymatically equivalent to those disclosed herein, wherein such DNA sequences are degenerate equivalents of the nucleic acid sequences disclosed herein in accordance with the degeneracy of the genetic code, is also encompassed by the present invention. Demonstration of the functionality of coding  
15 sequences identified by any of these methods can be carried out by complementation of mutants of appropriate organisms, such as *Synechocystis*, *Shewanella*, yeast, *Pseudomonas*, *Rhodobacteria*, etc., that lack specific biochemical reactions, or that have been mutated. The sequences of the DNA coding regions can be optimized by gene resynthesis, based on codon usage, for maximum expression in particular hosts.

20 For the alteration of tocopherol production in a host cell, a second expression construct can be used in accordance with the present invention. For example, the prenyltransferase expression construct can be introduced into a host cell in conjunction with a second expression construct having a nucleotide sequence for a protein involved in tocopherol biosynthesis.

25 The method of transformation in obtaining such transgenic plants is not critical to the instant invention, and various methods of plant transformation are currently available. Furthermore, as newer methods become available to transform crops, they may also be directly applied hereunder. For example, many plant species naturally susceptible to *Agrobacterium* infection may be successfully transformed via tripartite or binary vector  
30 methods of *Agrobacterium* mediated transformation. In many instances, it will be desirable to have the construct bordered on one or both sides by T-DNA, particularly having the left and right borders, more particularly the right border. This is particularly useful when the construct uses *A. tumefaciens* or *A. rhizogenes* as a mode for transformation, although the T-

DNA borders may find use with other modes of transformation. In addition, techniques of microinjection, DNA particle bombardment, and electroporation have been developed which allow for the transformation of various monocot and dicot plant species.

Normally, included with the DNA construct will be a structural gene having the  
5 necessary regulatory regions for expression in a host and providing for selection of transformant cells. The gene may provide for resistance to a cytotoxic agent, e.g. antibiotic, heavy metal, toxin, etc., complementation providing prototrophy to an auxotrophic host, viral immunity or the like. Depending upon the number of different host species the expression construct or components thereof are introduced, one or more markers may be employed,  
10 where different conditions for selection are used for the different hosts.

Where *Agrobacterium* is used for plant cell transformation, a vector may be used which may be introduced into the *Agrobacterium* host for homologous recombination with T-DNA or the Ti- or Ri-plasmid present in the *Agrobacterium* host. The Ti- or Ri-plasmid containing the T-DNA for recombination may be armed (capable of causing gall formation)  
15 or disarmed (incapable of causing gall formation), the latter being permissible, so long as the *vir* genes are present in the transformed *Agrobacterium* host. The armed plasmid can give a mixture of normal plant cells and gall.

In some instances where *Agrobacterium* is used as the vehicle for transforming host plant cells, the expression or transcription construct bordered by the T-DNA border region(s)  
20 will be inserted into a broad host range vector capable of replication in *E. coli* and *Agrobacterium*, there being broad host range vectors described in the literature. Commonly used is pRK2 or derivatives thereof. See, for example, Ditta, *et al.*, (*Proc. Nat. Acad. Sci., U.S.A.* (1980) 77:7347-7351) and EPA 0 120 515, which are incorporated herein by reference. Alternatively, one may insert the sequences to be expressed in plant cells into a vector  
25 containing separate replication sequences, one of which stabilizes the vector in *E. coli*, and the other in *Agrobacterium*. See, for example, McBride, *et al.* (*Plant Mol. Biol.* (1990) 14:269-276), wherein the pRiHRI (Jouanin, *et al.*, *Mol. Gen. Genet.* (1985) 201:370-374) origin of replication is utilized and provides for added stability of the plant expression vectors in host *Agrobacterium* cells.

30 Included with the expression construct and the T-DNA will be one or more markers, which allow for selection of transformed *Agrobacterium* and transformed plant cells. A number of markers have been developed for use with plant cells, such as resistance to chloramphenicol, kanamycin, the aminoglycoside G418, hygromycin, or the like. The

particular marker employed is not essential to this invention, one or another marker being preferred depending on the particular host and the manner of construction.

For transformation of plant cells using *Agrobacterium*, explants may be combined and incubated with the transformed *Agrobacterium* for sufficient time for transformation, the bacteria killed, and the plant cells cultured in an appropriate selective medium. Once callus forms, shoot formation can be encouraged by employing the appropriate plant hormones in accordance with known methods and the shoots transferred to rooting medium for regeneration of plants. The plants may then be grown to seed and the seed used to establish repetitive generations and for isolation of vegetable oils.

There are several possible ways to obtain the plant cells of this invention which contain multiple expression constructs. Any means for producing a plant comprising a construct having a DNA sequence encoding the expression construct of the present invention, and at least one other construct having another DNA sequence encoding an enzyme are encompassed by the present invention. For example, the expression construct can be used to transform a plant at the same time as the second construct either by inclusion of both expression constructs in a single transformation vector or by using separate vectors, each of which express desired genes. The second construct can be introduced into a plant which has already been transformed with the prenyltransferase expression construct, or alternatively, transformed plants, one expressing the prenyltransferase construct and one expressing the second construct, can be crossed to bring the constructs together in the same plant.

The nucleic acid sequences of the present invention can be used in constructs to provide for the expression of the sequence in a variety of host cells, both prokaryotic eukaryotic. Host cells of the present invention preferably include monocotyledenous and dicotyledenous plant cells.

In general, the skilled artisan is familiar with the standard resource materials which describe specific conditions and procedures for the construction, manipulation and isolation of macromolecules (e.g., DNA molecules, plasmids, etc.), generation of recombinant organisms and the screening and isolating of clones, (see for example, Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press (1989); Maliga *et al.*, *Methods in Plant Molecular Biology*, Cold Spring Harbor Press (1995), the entirety of which is herein incorporated by reference; Birren *et al.*, *Genome Analysis: Analyzing DNA*, 1, Cold Spring Harbor, New York, the entirety of which is herein incorporated by reference).



Methods for the expression of sequences in insect host cells are known in the art. Baculovirus expression vectors are recombinant insect viruses in which the coding sequence for a chosen foreign gene has been inserted behind a baculovirus promoter in place of the viral gene, e.g., polyhedrin (Smith and Summers, U.S. Pat. No., 4,745,051, the entirety of which is incorporated herein by reference). Baculovirus expression vectors are known in the art, and are described for example in Doerfler, *Curr. Top. Microbiol. Immunol.* 131:51-68 (1968); Luckow and Summers, *Bio/Technology* 6:47-55 (1988a); Miller, *Annual Review of Microbiol.* 42:177-199 (1988); Summers, *Curr. Comm. Molecular Biology*, Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (1988); Summers and Smith, *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*, Texas Ag. Exper. Station Bulletin No. 1555 (1988), the entireties of which is herein incorporated by reference)

Methods for the expression of a nucleic acid sequence of interest in a fungal host cell are known in the art. The fungal host cell may, for example, be a yeast cell or a filamentous fungal cell. Methods for the expression of DNA sequences of interest in yeast cells are generally described in "Guide to yeast genetics and molecular biology", Guthrie and Fink, eds. *Methods in enzymology*, Academic Press, Inc. Vol 194 (1991) and *Gene expression technology*, Goeddel ed, *Methods in Enzymology*, Academic Press, Inc., Vol 185 (1991).

Mammalian cell lines available as hosts for expression are known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC, Manassas, VA), such as HeLa cells, Chinese hamster ovary (CHO) cells, baby hamster kidney (BHK) cells and a number of other cell lines. Suitable promoters for mammalian cells are also known in the art and include, but are not limited to, viral promoters such as that from Simian Virus 40 (SV40) (Fiers *et al.*, *Nature* 273:113 (1978), the entirety of which is herein incorporated by reference), Rous sarcoma virus (RSV), adenovirus (ADV) and bovine papilloma virus (BPV). Mammalian cells may also require terminator sequences and poly-A addition sequences. Enhancer sequences which increase expression may also be included and sequences which promote amplification of the gene may also be desirable (for example methotrexate resistance genes).

Vectors suitable for replication in mammalian cells are well known in the art, and may include viral replicons, or sequences which insure integration of the appropriate sequences encoding epitopes into the host genome. Plasmid vectors that greatly facilitate the construction of recombinant viruses have been described (see, for example, Mackett *et al.*, *J Virol.* 49:857 (1984); Chakrabarti *et al.*, *Mol. Cell. Biol.* 5:3403 (1985); Moss, In: *Gene*

*Transfer Vectors For Mammalian Cells* (Miller and Calos, eds., Cold Spring Harbor Laboratory, N.Y., p. 10, (1987); all of which are herein incorporated by reference in their entirety).

The invention now being generally described, it will be more readily understood by reference to the following examples which are included for purposes of illustration only and are not intended to limit the present invention.

## EXAMPLES

### Example 1: Identification of Prenyltransferase Sequences

PSI-BLAST (Altschul, *et al.* (1997) *Nuc Acid Res* 25:3389-3402) profiles were generated for both the straight chain and aromatic classes of prenyltransferases. To generate the straight chain profile, a prenyl-transferase from *Porphyra purpurea* (Genbank accession 1709766) was used as a query against the NCBI non-redundant protein database. The *E. coli* enzyme involved in the formation of ubiquinone, *ubiA* (genbank accession 1790473) was used as a starting sequence to generate the aromatic prenyltransferase profile. These profiles were used to search public and proprietary DNA and protein data bases. In *Arabidopsis* seven putative prenyltransferases of the straight-chain class were identified, ATPT1, (SEQ ID NO:9), ATPT7 (SEQ ID NO:10), ATPT8 (SEQ ID NO:11), ATPT9 (SEQ ID NO:13), ATPT10 (SEQ ID NO:14), ATPT11 (SEQ ID NO:15), and ATPT12 (SEQ ID NO:16) and five were identified of the aromatic class, ATPT2 (SEQ ID NO:1), ATPT3 (SEQ ID NO:3), ATPT4 (SEQ ID NO:5), ATPT5 (SEQ ID NO:7), ATPT6 (SEQ ID NO:8). Additional prenyltransferase sequences from other plants related to the aromatic class of prenyltransferases, such as soy (SEQ ID NOs: 19-23, the deduced amino acid sequence of SEQ ID NO:23 is provided in SEQ ID NO:24) and maize (SEQ ID NOs:25-29, and 31) are also identified. The deduced amino acid sequence of ZMPT5 (SEQ ID NO:29) is provided in SEQ ID NO:30.

Searches are performed on a Silicon Graphics Unix computer using additional Bioaccelerator hardware and GenWeb software supplied by Compugen Ltd. This software and hardware enables the use of the Smith-Waterman algorithm in searching DNA and protein databases using profiles as queries. The program used to query protein databases is



profilesearch. This is a search where the query is not a single sequence but a profile based on a multiple alignment of amino acid or nucleic acid sequences. The profile is used to query a sequence data set, i.e., a sequence database. The profile contains all the pertinent information for scoring each position in a sequence, in effect replacing the "scoring matrix" used for the standard query searches. The program used to query nucleotide databases with a protein profile is tprofilesearch. Tprofilesearch searches nucleic acid databases using an amino acid profile query. As the search is running, sequences in the database are translated to amino acid sequences in six reading frames. The output file for tprofilesearch is identical to the output file for profilesearch except for an additional column that indicates the frame in which the best alignment occurred.

The Smith-Waterman algorithm, (Smith and Waterman (1981) *supra*), is used to search for similarities between one sequence from the query and a group of sequences contained in the database. E score values as well as other sequence information, such as conserved peptide sequences are used to identify related sequences.

To obtain the entire coding region corresponding to the *Arabidopsis* prenyltransferase sequences, synthetic oligo-nucleotide primers are designed to amplify the 5' and 3' ends of partial cDNA clones containing prenyltransferase sequences. Primers are designed according to the respective *Arabidopsis* prenyltransferase sequences and used in Rapid Amplification of cDNA Ends (RACE) reactions (Frohman *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:8998-9002) using the Marathon cDNA amplification kit (Clontech Laboratories Inc, Palo Alto, CA).

Additional BLAST searches are performed using the ATPT2 sequence, a sequence in the class of aromatic prenyl transferases. Additional sequences are identified in soybean libraries that are similar to the ATPT2 sequence. The additional soybean sequence demonstrates 80% identity and 91% similarity at the amino acid sequence.

Amino acid sequence alignments between ATPT2 (SEQ ID NO:2), ATPT3 (SEQ ID NO:4), ATPT4 (SEQ ID NO:6), ATPT8 (SEQ ID NO:12), and ATPT12 (SEQ ID NO:17) are performed using ClustalW (Figure 1), and the percent identity and similarities are provided in Table 1 below.

**Table 1:**

ATPT2	ATPT3	ATPT4	ATPT8	ATPT12
-------	-------	-------	-------	--------

ATPT2	% Identity		12	13	11	15
	% similar		25	25	22	32
	% Gap		17	20	20	9
ATPT3	% Identity			12	6	22
	% similar			29	16	38
	% Gap			20	24	14
ATPT4	% Identity				9	14
	% similar				18	29
	% Gap				26	19
ATPT8	% Identity					7
	% similar					19
	% Gap					20
ATPT12	% Identity					
	% similar					
	% Gap					

### Example 2: Preparation of Expression Constructs

5 A plasmid containing the napin cassette derived from pCGN3223 (described in USPN 5,639,790, the entirety of which is incorporated herein by reference) was modified to make it more useful for cloning large DNA fragments containing multiple restriction sites, and to allow the cloning of multiple napin fusion genes into plant binary transformation vectors. An adapter comprised of the self annealed oligonucleotide of sequence

10 CGCGATTAAATGGCGCGCCCTGCAGGCGGCCGCGCTGCAGGGCGCGCCATTAAAT (SEQ ID NO:40) was ligated into the cloning vector pBC SK+ (Stratagene) after digestion with the restriction endonuclease BssHII to construct vector pCGN7765. Plasmids pCGN3223 and pCGN7765 were digested with NotI and ligated together. The resultant vector, pCGN7770, contains the pCGN7765 backbone with the napin seed specific

15 expression cassette from pCGN3223.

The cloning cassette, pCGN7787, essentially the same regulatory elements as pCGN7770, with the exception of the napin regulatory regions of pCGN7770 have been

replaced with the double CAMV 35S promoter and the tml polyadenylation and transcriptional termination region.

A binary vector for plant transformation, pCGN5139, was constructed from pCGN1558 (McBride and Summerfelt, (1990) Plant Molecular Biology, 14:269-276). The polylinker of pCGN1558 was replaced as a HindIII/Asp718 fragment with a polylinker containing unique restriction endonuclease sites, AscI, PacI, XbaI, SwaI, BamHI, and NotI. The Asp718 and HindIII restriction endonuclease sites are retained in pCGN5139.

A series of turbo binary vectors are constructed to allow for the rapid cloning of DNA sequences into binary vectors containing transcriptional initiation regions (promoters) and transcriptional termination regions.

The plasmid pCGN8618 was constructed by ligating oligonucleotides 5'-TCGAGGATCCGCGGCCGCAAGCTTCCTGCAGG-3' (SEQ ID NO:41) and 5'-TCGACCTGCAGGAAGCTTGCGGCCGCGGATCC-3' (SEQ ID NO:42) into SalI/XhoI-digested pCGN7770. A fragment containing the napin promoter, polylinker and napin 3' region was excised from pCGN8618 by digestion with Asp718I; the fragment was blunt-ended by filling in the 5' overhangs with Klenow fragment then ligated into pCGN5139 that had been digested with Asp718I and HindIII and blunt-ended by filling in the 5' overhangs with Klenow fragment. A plasmid containing the insert oriented so that the napin promoter was closest to the blunted Asp718I site of pCGN5139 and the napin 3' was closest to the blunted HindIII site was subjected to sequence analysis to confirm both the insert orientation and the integrity of cloning junctions. The resulting plasmid was designated pCGN8622.

The plasmid pCGN8619 was constructed by ligating oligonucleotides 5'-TCGACCTGCAGGAAGCTTGCGGCCGCGGATCC-3' (SEQ ID NO:43) and 5'-TCGAGGATCCGCGGCCGCAAGCTTCCTGCAGG-3' (SEQ ID NO:44) into SalI/XhoI-digested pCGN7770. A fragment containing the napin promoter, polylinker and napin 3' region was removed from pCGN8619 by digestion with Asp718I; the fragment was blunt-ended by filling in the 5' overhangs with Klenow fragment then ligated into pCGN5139 that had been digested with Asp718I and HindIII and blunt-ended by filling in the 5' overhangs with Klenow fragment. A plasmid containing the insert oriented so that the napin promoter was closest to the blunted Asp718I site of pCGN5139 and the napin 3' was closest to the blunted HindIII site was subjected to sequence analysis to confirm both the insert orientation and the integrity of cloning junctions. The resulting plasmid was designated pCGN8623.

The plasmid pCGN8620 was constructed by ligating oligonucleotides 5'-TCGAGGATCCGCGGCCGCAAGCTTCCTGCAGGAGCT -3' (SEQ ID NO:45) and 5'-CCTGCAGGAAGCTTGCGGCCGCGGATCC-3' (SEQ ID NO:46) into SalI/SacI-digested pCGN7787. A fragment containing the d35S promoter, polylinker and tml 3' region was removed from pCGN8620 by complete digestion with Asp718I and partial digestion with NotI. The fragment was blunt-ended by filling in the 5' overhangs with Klenow fragment then ligated into pCGN5139 that had been digested with Asp718I and HindIII and blunt-ended by filling in the 5' overhangs with Klenow fragment. A plasmid containing the insert oriented so that the d35S promoter was closest to the blunted Asp718I site of pCGN5139 and the tml 3' was closest to the blunted HindIII site was subjected to sequence analysis to confirm both the insert orientation and the integrity of cloning junctions. The resulting plasmid was designated pCGN8624.

The plasmid pCGN8621 was constructed by ligating oligonucleotides 5'-TCGACCTGCAGGAAGCTTGCGGCCGCGGATCCAGCT -3' (SEQ ID NO:47) and 5'-GGATCCGCGGCCGCAAGCTTCCTGCAGG-3' (SEQ ID NO:48) into SalI/SacI-digested pCGN7787. A fragment containing the d35S promoter, polylinker and tml 3' region was removed from pCGN8621 by complete digestion with Asp718I and partial digestion with NotI. The fragment was blunt-ended by filling in the 5' overhangs with Klenow fragment then ligated into pCGN5139 that had been digested with Asp718I and HindIII and blunt-ended by filling in the 5' overhangs with Klenow fragment. A plasmid containing the insert oriented so that the d35S promoter was closest to the blunted Asp718I site of pCGN5139 and the tml 3' was closest to the blunted HindIII site was subjected to sequence analysis to confirm both the insert orientation and the integrity of cloning junctions. The resulting plasmid was designated pCGN8625.

The plasmid construct pCGN8640 is a modification of pCGN8624 described above. A 938bp PstI fragment isolated from transposon Tn7 which encodes bacterial spectinomycin and streptomycin resistance (Fling et al. (1985), *Nucleic Acids Research* 13(19):7095-7106), a determinant for E. coli and Agrobacterium selection, was blunt ended with Pfu polymerase. The blunt ended fragment was ligated into pCGN8624 that had been digested with SpeI and blunt ended with Pfu polymerase. The region containing the PstI fragment was sequenced to confirm both the insert orientation and the integrity of cloning junctions.

The spectinomycin resistance marker was introduced into pCGN8622 and pCGN8623 as follows. A 7.7 Kbp AvrII-SnaBI fragment from pCGN8640 was ligated to a 10.9 Kbp

AvrII-SnaBI fragment from pCGN8623 or pCGN8622, described above. The resulting plasmids were pCGN8641 and pCGN8643, respectively.

The plasmid pCGN8644 was constructed by ligating oligonucleotides 5'-GATCACCTGCAGGAAGCTTGCGGCCGCGGATCCAATGCA-3' (SEQ ID NO:49) and 5'-TTGGATCCGCGGCCGCAAGCTTCCTGCAGGT-3' (SEQ ID NO:50) into BamHI-PstI digested pCGN8640.

Synthetic oligonucleotides were designed for use in Polymerase Chain Reactions (PCR) to amplify the coding sequences of ATPT2, ATPT3, ATPT4, ATPT8, and ATPT12 for the preparation of expression constructs and are provided in Table 2 below.

**Table 2:**

Name	Restriction Site	Sequence	SEQ ID NO:
ATPT2	5' NotI	GGATCCGCGGCCGCGCACAAATGGAGTC TCTGCTCTCTAGTTCT	51
ATPT2	3' SseI	GGATCCTGCAGGTCACCTCAAAAAA GGTAACAGCAAGT	52
ATPT3	5' NotI	GGATCCGCGGCCGCGCACAAATGGCGTT TTTTGGGCTCTCCCGTGTTT	53
ATPT3	3' SseI	GGATCCTGCAGGTTATTGAAAACCTT CTCCAAGTACAAC	54
ATPT4	5' NotI	GGATCCGCGGCCGCGCACAAATGTGGCG AAGATCTGTTGTT	55
ATPT4	3' SseI	GGATCCTGCAGGTCATGGAGAGTAG AAGGAAGGAGCT	56
ATPT8	5' NotI	GGATCCGCGGCCGCGCACAAATGGTACT TGCCGAGGTTCCAAAGCTTGCCTCT	57
ATPT8	3' SseI	GGATCCTGCAGGTCACCTGTTTCTG GTGATGACTCTAT	58
ATPT12	5' NotI	GGATCCGCGGCCGCGCACAAATGACTTC GATTCTCAACACT	59
ATPT12	3' SseI	GGATCCTGCAGGTCAGTGTTGCGAT GCTAATGCCGT	60

The coding sequences of ATPT2, ATPT3, ATPT4, ATPT8, and ATPT12 were all amplified using the respective PCR primers shown in Table 2 above and cloned into the TopoTA vector (Invitrogen). Constructs containing the respective prenyltransferase sequences were digested with NotI and Sse8387I and cloned into the turbobinary vectors described above.

The sequence encoding ATPT2 prenyltransferase was cloned in the sense orientation into pCGN8640 to produce the plant transformation construct pCGN10800 (Figure 2). The ATPT2 sequence is under control of the 35S promoter.

The ATPT2 sequence was also cloned in the antisense orientation into the construct pCGN8641 to create pCGN10801 (Figure 3). This construct provides for the antisense expression of the ATPT2 sequence from the napin promoter.

5 The ATPT2 coding sequence was also cloned in the antisense orientation into the vector pCGN8643 to create the plant transformation construct pCGN10802

The ATPT2 coding sequence was also cloned in the antisense orientation into the vector pCGN8644 to create the plant transformation construct pCGN10803 (Figure 4).

10 The ATPT4 coding sequence was cloned into the vector pCGN864 to create the plant transformation construct pCGN10806 (Figure 5). The ATPT2 coding sequence was cloned into the vector pCGN864 to create the plant transformation construct pCGN10807 (Figure 6). The ATPT3 coding sequence was cloned into the vector pCGN864 to create the plant transformation construct pCGN10808 (Figure 7). The ATPT3 coding sequence was cloned in the sense orientation into the vector pCGN8640 to create the plant transformation construct pCGN10809 (Figure 8). The ATPT3 coding sequence was cloned in the antisense orientation into the vector  
15 pCGN8641 to create the plant transformation construct pCGN10810 (Figure 9). The ATPT3 coding sequence was cloned into the vector pCGN8643 to create the plant transformation construct pCGN10811 (Figure 10). The ATPT3 coding sequence was cloned into the vector pCGN8640 to create the plant transformation construct pCGN10812 (Figure 11). The ATPT4 coding sequence was cloned into the vector pCGN8640 to create the plant transformation  
20 construct pCGN10813 (Figure 12). The ATPT4 coding sequence was cloned into the vector pCGN8643 to create the plant transformation construct pCGN10814 (Figure 13). The ATPT4 coding sequence was cloned into the vector pCGN8641 to create the plant transformation construct pCGN10815 (Figure 14). The ATPT4 coding sequence was cloned in the antisense orientation into the vector pCGN8644 to create the plant transformation construct pCGN10816  
25 (Figure 15). The ATPT2 coding sequence was cloned into the vector pCGN???? to create the plant transformation construct pCGN10817 (Figure 16). The ATPT8 coding sequence was cloned in the sense orientation into the vector pCGN8643 to create the plant transformation construct pCGN10819 (Figure 17). The ATPT12 coding sequence was cloned into the vector pCGN8644 to create the plant transformation construct pCGN10824 (Figure 18). The ATPT12 coding  
30 sequence was cloned into the vector pCGN8641 to create the plant transformation construct pCGN10825 (Figure 19). The ATPT8 coding sequence was cloned into the vector pCGN8644 to create the plant transformation construct pCGN10826 (Figure 20).

**Example 3: Plant Transformation**

5 Transgenic *Brassica* plants are obtained by *Agrobacterium*-mediated transformation as described by Radke *et al.* (*Theor. Appl. Genet.* (1988) 75:685-694; *Plant Cell Reports* (1992) 11:499-505). Transgenic *Arabidopsis thaliana* plants may be obtained by *Agrobacterium*-mediated transformation as described by Valverkens *et al.*, (*Proc. Nat. Acad. Sci.* (1988) 85:5536-5540), or as described by Bent *et al.* ((1994), *Science* 265:1856-1860), or  
 10 Bechtold *et al.* ((1993), *C.R.Acad.Sci, Life Sciences* 316:1194-1199). Other plant species may be similarly transformed using related techniques.

Alternatively, microprojectile bombardment methods, such as described by Klein *et al.* (*Bio/Technology* 10:286-291) may also be used to obtain nuclear transformed plants.

15

**Example 4: Identification of Additional Prenyltransferases**

A PSI-Blast profile generated using the *E. coli* *ubiA* (genbank accession 1790473) sequence was used to analyze the *Synechocystis* genome. This analysis identified 5 open  
 20 reading frames (ORFs) in the *Synechocystis* genome that were potentially prenyltransferases; *slr0926* (annotated as *ubiA* (4-hydroxybenzoate-octaprenyl transferase, SEQ ID NO:32), *slr1899* (annotated as *ctaB* (cytochrome c oxidase folding protein, SEQ ID NO:33), *slr0056* (annotated as *g4* (chlorophyll synthase 33 kd subunit, SEQ ID NO:34), *slr1518* (annotated as *menA* (menaquinone biosynthesis protein, SEQ ID NO:35), and *slr1736* (annotated as a  
 25 hypothetical protein of unknown function (SEQ ID NO:36).

To determine the functionality of these ORFs and their involvement, if any, in the biosynthesis of Tocopherols, knockouts constructs were made to disrupt the ORF identified in *Synechocystis*.

Synthetic oligos were designed to amplify regions from the 5' (5'-  
 30 TAATGTGTACATTGTCGGCCTC (17365') (SEQ ID NO:61) and 5'-  
 GCAATGTAACATCAGAGATTTTGAGACACAACGTGGCTTTCCACAATTCCCCGCA  
 CCGTC (1736kanpr1)) (SEQ ID NO:62) and 3' (5'-AGGCTAATAAGCACAAATGGGA  
 (17363') (SEQ ID NO:63) and 5'-



GGTATGAGTCAGCAACACCTTCTTCACGAGGCAGACCTCAGC

GGAATTGGTTTATAGGTTATCCC (1736kanpr2)) (SEQ ID NO:64) ends of the slr1736 ORF.

The 1736kanpr1 and 1736kanpr2 oligos contained 20 bp of homology to the slr1736 ORF with an additional 40 bp of sequence homology to the ends of the kanamycin resistance

5 cassette. Separate PCR steps were completed with these oligos and the products were gel purified and combined with the kanamycin resistance gene from puc4K (Pharmacia) that had been digested with *HincII* and gel purified away from the vector backbone. The combined fragments were allowed to assemble without oligos under the following conditions: 94°C for 1 min, 55°C for 1 min, 72°C for 1 min plus 5 seconds per cycle for 40 cycles using pfu  
10 polymerase in 100ul reaction volume (Zhao, H and Arnold (1997) *Nucleic Acids Res.* 25(6):1307-1308). One microliter or five microliters of this assembly reaction was then amplified using 5' and 3' oligos nested within the ends of the ORF fragment, so that the resulting product contained 100-200 bp of the 5' end of the *Synechocystis* gene to be knocked out, the kanamycin resistance cassette, and 100-200 bp of the 3' end of the gene to be  
15 knocked out. This PCR product was then cloned into the vector pGemT easy (Promega) to create the construct pMON21681 and used for *Synechocystis* transformation.

Primers were also synthesized for the preparation of *Synechocystis* knockout constructs for the other sequences using the same method as described above, with the following primers. The *ubiA* 5' sequence was amplified using the primers 5'-

20 GGATCCATGGTT GCCCAAACCCCATC (SEQ ID NO:65) and 5'-

GCAATGTAACATCAGAGA TTTTGAGACACAACG

TGGCTTTGGGTAAGCAACAATGACCGGC (SEQ ID NO:66). The 3' region was amplified using the synthetic oligonucleotide primers 5'-

GAATTCTCAAAGCCAGCCCAGTAAC (SEQ ID NO:67) and 5'-GGTATGAGTC

25 AGCAACACCTTCTTCACGAGGCAGACCTCAGCGGGTGCGAAAAGGGTTTTCCC

(SEQ ID NO:68). The amplification products were combined with the kanamycin resistance gene from puc4K (Pharmacia) that had been digested with *HincII* and gel purified away from the vector backbone. The annealed fragment was amplified using 5' and 3' oligos nested

within the ends of the ORF fragment (5'-CCAGTGGTTTATAGGCTGTGTGGTC (SEQ ID

30 NO:69) and 5'-CTGAGTTGGATGTATTGGATC (SEQ ID NO:70)), so that the resulting

product contained 100-200 bp of the 5' end of the *Synechocystis* gene to be knocked out, the kanamycin resistance cassette, and 100-200 bp of the 3' end of the gene to be knocked out.



This PCR product was then cloned into the vector pGemT easy (Promega) to create the construct pMON21682 and used for *Synechocystis* transformation.

Primers were also synthesized for the preparation of *Synechocystis* knockout constructs for the other sequences using the same method as described above, with the

5 following primers. The sl11899 5' sequence was amplified using the primers 5'-

GGATCCATGGTTACTT CGACAAAAATCC (SEQ ID NO:71) and 5'-

GCAATGTAACATCAGAG

ATTTTGAGACACAACGTGGCTTTGCTAGGCAACCGCTTAGTAC (SEQ ID NO:72).

The 3' region was amplified using the synthetic oligonucleotide primers 5'-

10 GAATTCTTAACCCAACAGTAAAGTTCCC (SEQ ID NO:73) and 5'-

GGTATGAGTCAGC

AACACCTTCTTCACGAGGCAGACCTCAGCGCCGGCATTGTCTTTTACATG (SEQ ID

NO:74). The amplification products were combined with the kanamycin resistance gene from puc4K (Pharmacia) that had been digested with *HincII* and gel purified away from the vector

15 backbone. The annealed fragment was amplified using 5' and 3' oligos nested within the

ends of the ORF fragment (5'- GGAACCCTTGCAGCCGCTTC (SEQ ID NO:75)

and 5'- GTATGCCCAACTGGTGCAGAGG (SEQ ID NO:76)), so that the resulting product

contained 100-200 bp of the 5' end of the *Synechocystis* gene to be knocked out, the

kanamycin resistance cassette, and 100-200 bp of the 3' end of the gene to be knocked out.

20 This PCR product was then cloned into the vector pGemT easy (Promega) to create the

construct pMON21679 and used for *Synechocystis* transformation.

Primers were also synthesized for the preparation of *Synechocystis* knockout

constructs for the other sequences using the same method as described above, with the

following primers. The slr0056 5' sequence was amplified using the primers 5'-

25 GGATCCATGTCTGACACACAAAATACCG (SEQ ID NO:77) and 5'-

GCAATGTAACATCAGAGATTTTGAGACACAACGTGGCTTTCGCCAATACCAGCCA

CCAACAG (SEQ ID NO:78). The 3' region was amplified using the synthetic

oligonucleotide primers 5'- GAATTCTCAAAT CCCCGCATGGCCTAG (SEQ ID NO:79)

and 5'-

30 GGTATGAGTCAGCAACACCTTCTTCACGAGGCAGACCTCAGCGGCCTACGGCTTG

GACGTGTGGG (SEQ ID NO:80). The amplification products were combined with the

kanamycin resistance gene from puc4K (Pharmacia) that had been digested with *HincII* and

gel purified away from the vector backbone. The annealed fragment was amplified using 5'

and 3' oligos nested within the ends of the ORF fragment (5'-

CACTTGGATTCCCCTGATCTG (SEQ ID NO:81) and 5'-

GCAATACCCGCTTGGAAAACG (SEQ ID NO:82)), so that the resulting product

contained 100-200 bp of the 5' end of the *Synechocystis* gene to be knocked out, the

5 kanamycin resistance cassette, and 100-200 bp of the 3' end of the gene to be knocked out.

This PCR product was then cloned into the vector pGemT easy (Promega) to create the construct pMON21677 and used for *Synechocystis* transformation.

Primers were also synthesized for the preparation of *Synechocystis* knockout constructs for the other sequences using the same method as described above, with the

10 following primers. The slr1518 5' sequence was amplified using the primers 5'-

GGATCCATGACCGAAT CTTCGCCCCTAGC (SEQ ID NO:83) and 5'-

GCAATGTAACATCAGAGATTTTGA GACACAACGTGGC

TTTCAATCCTAGGTAGCCGAGGCG (SEQ ID NO:84). The 3' region was amplified

using the synthetic oligonucleotide primers 5'- GAATTCTTAGCCCAGGCC AGCCCAGCC

15 (SEQ ID NO:85) and 5'- GGTATGAGTCAGCAACACCTTCTTCACGA

GGCAGACCTCAGCGGGGAATTGATTTGTTTAATTACC (SEQ ID NO:86). The

amplification products were combined with the kanamycin resistance gene from puc4K

(Pharmacia) that had been digested with *HincII* and gel purified away from the vector

backbone. The annealed fragment was amplified using 5' and 3' oligos nested within the

20 ends of the ORF fragment (5'- GCGATCGCCATTATCGCTTGG (SEQ ID NO:87) and 5'-

GCAGACTGGCAATTATCAGTAACG (SEQ ID NO:88)), so that the resulting product

contained 100-200 bp of the 5' end of the *Synechocystis* gene to be knocked out, the

kanamycin resistance cassette, and 100-200 bp of the 3' end of the gene to be knocked out.

This PCR product was then cloned into the vector pGemT easy (Promega) to create the

25 construct pMON21680 and used for *Synechocystis* transformation.

## B. Transformation of *Synechocystis*

Cells of *Synechocystis* 6803 were grown to a density of approximately  $2 \times 10^8$  cells per ml and harvested by centrifugation. The cell pellet was re-suspended in fresh BG-11 medium

30 (ATCC Medium 616) at a density of  $1 \times 10^9$  cells per ml and used immediately for

transformation. One-hundred microliters of these cells were mixed with 5 ul of mini prep

DNA and incubated with light at 30C for 4 hours. This mixture was then plated onto nylon

filters resting on BG-11 agar supplemented with TES pH8 and allowed to grow for 12-18

hours. The filters were then transferred to BG-11 agar + TES + 5ug/ml kanamycin and allowed to grow until colonies appeared within 7-10 days (Packer and Glazer, 1988). Colonies were then picked into BG-11 liquid media containing 5 ug/ml kanamycin and allowed to grow for 5 days. These cells were then transferred to Bg-11 media containing  
 5 10ug/ml kanamycin and allowed to grow for 5 days and then transferred to Bg-11 + kanamycin at 25ug/ml and allowed to grow for 5 days. Cells were then harvested for PCR analysis to determine the presence of a disrupted ORF and also for HPLC analysis to determine if the disruption had any effect on tocopherol levels.

PCR analysis of the *Synechocystis* isolates for slr1736 and sll1899 showed complete  
 10 segregation of the mutant genome, meaning no copies of the wild type genome could be detected in these strains. This suggests that function of the native gene is not essential for cell function. HPLC analysis of these same isolates showed that the sll1899 strain had no detectable reduction in tocopherol levels. However, the strain carrying the knockout for slr1736 produced no detectable levels of tocopherol.

15 The amino acid sequences for the *Synechocystis* knockouts are compared using ClustalW, and are provided in Table 3 below. Provided are the percent identities, percent similarity, and the percent gap. The alignment of the sequences is provided in Figure 21.

**Table 3:**

	Slr1736	slr0926	sll1899	slr0056	slr1518
slr1736 %identity		14	12	18	11
%similar		29	30	34	26
%gap		8	7	10	5
slr0926 %identity			20	19	14
%similar			39	32	28
%gap			7	9	4
sll1899 %identity				17	13
%similar				29	29
%gap				12	9
slr0056 %identity					15
%similar					31
%gap					8

slr1518 %identity	
%similar	
%gap	

Amino acid sequence comparisons are performed using various *Arabidopsis* prenyltransferase sequences and the *Synechocystis* sequences. The comparisons are presented in Table 4 below. Provided are the percent identities, percent similarity, and the percent gap.

5 The alignment of the sequences is provided in Figure 22.

**Table 4:**

	ATPT2	slr1736	ATPT3	slr0926	ATPT4	sl11899	ATPT12	slr0056	ATPT8	slr1518
ATPT2		29	9	9	8	8	12	9	7	9
		46	23	21	20	20	28	23	21	20
		27	13	28	23	29	11	24	25	24
slr1736			9	13	8	12	13	15	8	10
			19	28	19	28	26	33	21	26
			34	12	34	15	26	10	12	10
ATPT3				23	11	14	13	10	5	11
				36	26	26	26	21	14	22
				29	21	31	16	30	30	30
					12	20	17	20	11	14
slr0926					24	37	28	33	24	29
					33	12	25	10	11	9
						18	11	8	6	7
ATPT4						33	23	18	16	19
						28	19	32	32	33
							13	17	10	12
sl11899							24	30	23	26
							27	13	10	11
								52	8	11
ATPT1								66	19	26
2										
								18	25	23

		9	13
slr0056		23	32
		10	8
			7
ATPT8			23
			7
slr1518			

#### 4B. Preparation of the slr1737 Knockout

The *Synechocystis* sp. 6803 slr1737 knockout was constructed by the following method. The GPS™-1 Genome Priming System (New England Biolabs) was used to insert, by a Tn7 Transposase system, a Kanamycin resistance cassette into *slr1737*. A plasmid from a *Synechocystis* genomic library clone containing 652 base pairs of the targeted orf (*Synechocystis* genome base pairs 1324051 – 1324703; the predicted orf base pairs 1323672 – 1324763, as annotated by Cyanobase) was used as target DNA. The reaction was performed according to the manufacturers protocol. The reaction mixture was then transformed into *E. coli* DH10B electrocompetant cells and plated. Colonies from this transformation were then screened for transposon insertions into the target sequence by amplifying with M13 Forward and Reverse Universal primers, yielding a product of 652 base pairs plus ~1700 base pairs, the size of the transposon kanamycin cassette, for a total fragment size of ~2300 base pairs. After this determination, it was then necessary to determine the approximate location of the insertion within the targeted orf, as 100 base pairs of orf sequence was estimated as necessary for efficient homologous recombination in *Synechocystis*. This was accomplished through amplification reactions using either of the primers to the ends of the transposon, Primer S (5' end) or N (3' end), in combination with either a M13 Forward or Reverse primer. That is, four different primer combinations were used to map each potential knockout construct: Primer S – M13 Forward, Primer S – M13 Reverse, Primer N – M13 Forward, Primer N – M13 Reverse. The construct used to transform *Synechocystis* and knockout slr1737 was determined to consist of a approximately

150 base pairs of slr1737 sequence on the 5' side of the transposon insertion and approximately 500 base pairs on the 3' side, with the transcription of the orf and kanamycin cassette in the same direction. The nucleic acid sequence of slr1737 is provided in SEQ ID NO:38 the deduced amino acid sequence is provided in SEQ ID NO:39.

5 Cells of *Synechocystis* 6803 were grown to a density of  $\sim 2 \times 10^8$  cells per ml and harvested by centrifugation. The cell pellet was re-suspended in fresh BG-11 medium at a density of  $1 \times 10^9$  cells per ml and used immediately for transformation. 100  $\mu$ l of these cells were mixed with 5  $\mu$ l of mini prep DNA and incubated with light at 30C for 4 hours. This mixture was then plated onto nylon filters resting on BG-11 agar supplemented with TES pH8  
10 and allowed to grow for 12-18 hours. The filters were then transferred to BG-11 agar + TES + 5  $\mu$ g/ml kanamycin and allowed to grow until colonies appeared within 7-10 days (Packer and Glazer, 1988). Colonies were then picked into BG-11 liquid media containing 5  $\mu$ g/ml kanamycin and allowed to grow for 5 days. These cells were then transferred to Bg-11 media containing 10  $\mu$ g/ml kanamycin and allowed to grow for 5 days and then transferred to Bg-11  
15 + kanamycin at 25  $\mu$ g/ml and allowed to grow for 5 days. Cells were then harvested for PCR analysis to determine the presence of a disrupted ORF and also for HPLC analysis to determine if the disruption had any effect on tocopherol levels.

PCR analysis of the *Synechocystis* isolates, using primers to the ends of the *slr1737* orf, showed complete segregation of the mutant genome, meaning no copies of the wild type  
20 genome could be detected in these strains. This suggests that function of the native gene is not essential for cell function. HPLC analysis of the strain carrying the knockout for *slr1737* produced no detectable levels of tocopherol.

#### 4C. Phytyl Prenyltransferase Enzyme Assays

25 [ $^3$ H] Homogentisic acid in 0.1%  $\text{H}_3\text{PO}_4$  (specific radioactivity 40 Ci/mmol). Phytyl pyrophosphate was synthesized as described by Joo, *et al.* (1973) *Can J. Biochem.* 51:1527. 2-methyl-6-phytylquinol and 2,3-dimethyl-5-phytylquinol were synthesized as described by Soll, *et al.* (1980) *Phytochemistry* 19:215. Homogentisic acid,  $\alpha$ ,  $\beta$ ,  $\delta$ , and  $\gamma$ -tocopherol, and tocol, were purchased commercially.

30 The wild-type strain of *Synechocystis* sp. PCC 6803 was grown in BG11 medium with bubbling air at 30°C under  $50 \mu\text{E} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$  fluorescent light, and 70% relative humidity. The growth medium of slr1736 knock-out (potential PPT) strain of this organism was

supplemented with  $25 \mu\text{g mL}^{-1}$  kanamycin. Cells were collected from 0.25 to 1 liter culture by centrifugation at 5000 g for 10 min and stored at  $-80^{\circ}\text{C}$ .

Total membranes were isolated according to Zak's procedures with some modifications (Zak, *et al.* (1999) *Eur J. Biochem* 261:311). Cells were broken on a French press. Before the  
5 French press treatment, the cells were incubated for 1 hour with lysozyme (0.5%, w/v) at  $30^{\circ}\text{C}$  in a medium containing 7 mM EDTA, 5 mM NaCl and 10 mM Hepes-NaOH, pH 7.4. The spheroplasts were collected by centrifugation at 5000 g for 10 min and resuspended at 0.1 - 0.5 mg chlorophyll $\cdot\text{mL}^{-1}$  in 20 mM potassium phosphate buffer, pH 7.8. Proper amount of protease inhibitor cocktail and DNAase I from Boehringer Mannheim were added to the  
10 solution. French press treatments were performed two to three times at 100 MPa. After breakage, the cell suspension was centrifuged for 10 min at 5000g to pellet unbroken cells, and this was followed by centrifugation at 100 000 g for 1 hour to collect total membranes. The final pellet was resuspended in a buffer containing 50 mM Tris-HCL and 4 mM  $\text{MgCl}_2$ .

Chloroplast pellets were isolated from 250 g of spinach leaves obtained from local  
15 markets. Devined leaf sections were cut into grinding buffer (2 l /250 g leaves) containing 2 mM EDTA, 1 mM  $\text{MgCl}_2$ , 1 mM  $\text{MnCl}_2$ , 0.33 M sorbitol, 0.1% ascorbic acid, and 50 mM Hepes at pH 7.5. The leaves were homogenized for 3 sec three times in a 1-L blender, and filtered through 4 layers of miracloth. The supernatant was then centrifuged at 5000g for 6 min. The chloroplast pellets were resuspended in small amount of grinding buffer (Douce,*et*  
20 *al* Methods in Chloroplast Molecular Biology, 239 (1982)

Chloroplasts in pellets can be broken in three ways. Chloroplast pellets were first aliquoted in 1 mg of chlorophyll per tube, centrifuged at 6000 rpm for 2 min in microcentrifuge, and grinding buffer was removed. Two hundred microliters of Triton X-100 buffer (0.1% Triton X-100, 50 mM Tris-HCl pH 7.6 and 4 mM  $\text{MgCl}_2$ ) or swelling buffer (10  
25 mM Tris pH 7.6 and 4 mM  $\text{MgCl}_2$ ) was added to each tube and incubated for  $\frac{1}{2}$  hour at  $4^{\circ}\text{C}$ . Then the broken chloroplast pellets were used for the assay immediately. In addition, broken chloroplasts can also be obtained by freezing in liquid nitrogen and stored at  $-80^{\circ}\text{C}$  for  $\frac{1}{2}$  hour, then used for the assay.

In some cases chloroplast pellets were further purified with 40%/ 80% percoll gradient to  
30 obtain intact chloroplasts. The intact chloroplasts were broken with swelling buffer, then either used for assay or further purified for envelope membranes with 20.5%/ 31.8% sucrose density gradient (Sol, *et al* (1980) *supra*). The membrane fractions were centrifuged at 100 000g for 40 min and resuspended in 50 mM Tris-HCl pH 7.6, 4 mM  $\text{MgCl}_2$ .



Various amounts of [ $^3\text{H}$ ]HGA, 40 to 60  $\mu\text{M}$  unlabelled HGA with specific activity in the range of 0.16 to 4 Ci/mmol were mixed with a proper amount of 1M Tris-NaOH pH 10 to adjust pH to 7.6. HGA was reduced for 4 min with a trace amount of solid  $\text{NaBH}_4$ . In addition to HGA, standard incubation mixture (final vol 1 mL) contained 50 mM Tris-HCl, pH 7.6, 3-5 mM  $\text{MgCl}_2$ , and 100  $\mu\text{M}$  phytyl pyrophosphate. The reaction was initiated by addition of *Synechocystis* total membranes, spinach chloroplast pellets, spinach broken chloroplasts, or spinach envelope membranes. The enzyme reaction was carried out for 2 hour at 23°C or 30°C in the dark or light. The reaction is stopped by freezing with liquid nitrogen, and stored at -80°C or directly by extraction.

A constant amount of tocol was added to each assay mixture and reaction products were extracted with a 2 mL mixture of chloroform/methanol (1:2, v/v) to give a monophasic solution. NaCl solution (2 mL; 0.9%) was added with vigorous shaking. This extraction procedure was repeated three times. The organic layer containing the prenylquinones was filtered through a 20  $\mu\text{m}$  filter, evaporated under  $\text{N}_2$ , and then resuspended in 100  $\mu\text{L}$  of ethanol.

The samples were mainly analyzed by Normal-Phase HPLC method (Isocratic 90% Hexane and 10% Methyl-t-butyl ether), and use a Zorbax silica column, 4.6 x 250 mm. The samples were also analyzed by Reversed-Phase HPLC method (Isocratic 0.1%  $\text{H}_3\text{PO}_4$  in MeOH), and use a Vydac 201HS54 C18 column; 4.6 x 250 mm coupled with an All-tech C18 guard column. The amount of products were calculated based on the substrate specific radioactivity, and adjusted according to the % recovery based on the amount of internal standard.

The amount of chlorophyll was determined as described in Arnon (1949) *Plant Physiol.* 24:1. Amount of protein was determined by the Bradford method using gamma globulin as a standard (Bradford, (1976) *Anal. Biochem.* 72:248)

Results of the assay demonstrate that 2-Methyl-6-Phytylplastoquinone is produced in the *Synechocystis* slr1736 knockout preparations. The results of the phytyl prenyltransferase enzyme activity assay for the slr1736 knock out are presented in Figure 23.

#### 4D. Complementation of the slr1736 knockout with ATPT2

In order to determine whether ATPT2 could complement the knockout of slr1736 in *Synechocystis* 6803 a plasmid was constructed to express the ATPT2 sequence from the TAC promoter. A vector, plasmid psl1211, was obtained from the lab of Dr. Himadri Pakrasi of



Washington University, and is based on the plasmid RSF1010 which is a broad host range plasmid (Ng W.-O., Zentella R., Wang, Y., Taylor J-S. A., Pakrasi, H.B. 2000. *phrA*, the major photoreactivating factor in the cyanobacterium *Synechocystis* sp. strain PCC 6803 codes for a cyclobutane pyrimidine dimer specific DNA photolyase. *Arch. Microbiol.* (in press)). The ATPT2 gene was isolated from the vector pCGN10817 by PCR using the following primers. ATPT2nco.pr 5'-CCATGGATTCGAGTAAAGTTGTCGC (SEQ ID NO:89); ATPT2ri.pr- 5'-GAATTCACCTTCAAAAAAGGTAACAG (SEQ ID NO:90). These primers will remove approximately 112 BP from the 5' end of the ATPT2 sequence, which is thought to be the chloroplast transit peptide. These primers will also add an NcoI site at the 5' end and an EcoRI site at the 3' end which can be used for sub-cloning into subsequent vectors. The PCR product from using these primers and pCGN10817 was ligated into pGEM T easy and the resulting vector pMON21689 was confirmed by sequencing using the m13forward and m13reverse primers. The NcoI/EcoRI fragment from pMON21689 was then ligated with the EagI/EcoRI and EagI/NcoI fragments from psl1211 resulting in pMON21690. The plasmid pMON21690 was introduced into the slr1736 *Synechocystis* 6803 KO strain via conjugation. Cells of sl906 (a helper strain) and DH10B cells containing pMON21690 were grown to log phase (O.D. 600= 0.4) and 1 ml was harvested by centrifugation. The cell pellets were washed twice with a sterile BG-11 solution and resuspended in 200 ul of BG-11. The following was mixed in a sterile eppendorf tube: 50 ul SL906, 50 ul DH10B cells containing pMON21690, and 100 ul of a fresh culture of the slr1736 *Synechocystis* 6803 KO strain (O.D. 730 = 0.2-0.4). The cell mixture was immediately transferred to a nitrocellulose filter resting on BG-11 and incubated for 24 hours at 30C and 2500 LUX(50 ue) of light. The filter was then transferred to BG-11 supplemented with 10ug/ml Gentamycin and incubated as above for ~5 days. When colonies appeared, they were picked and grown up in liquid BG-11 + Gentamycin 10 ug/ml. (Elhai, J. and Wolk, P. 1988. Conjugal transfer of DNA to Cyanobacteria. *Methods in Enzymology* 167, 747-54) The liquid cultures were then assayed for tocopherols by harvesting 1ml of culture by centrifugation, extracting with ethanol/pyrogallol, and HPLC separation. The slr1736 *Synechocystis* 6803 KO strain, did not contain any detectable tocopherols, while the slr1736 *Synechocystis* 6803 KO strain transformed with pmon21690 contained detectable alpha tocopherol. A *Synechocystis* 6803 strain transformed with psl1211(vector control) produced alpha tocopherol as well.

**Example 5: Transgenic Plant Analysis**

Arabidopsis plants transformed with constructs for the sense or antisense expression of the ATPT proteins were analyzed by High Pressure Liquid Chromatography (HPLC) for altered levels of total tocopherols, as well as altered levels of specific tocopherols (alpha, beta, gamma, and delta tocopherol).

Extracts of leaves and seeds were prepared for HPLC as follows. For seed extracts, 10 mg of seed was added to 1 g of microbeads (Biospec) in a sterile microfuge tube to which 500 ul 1% pyrogallol (Sigma Chem)/ethanol was added. The mixture was shaken for 3 minutes in a mini Beadbeater (Biospec) on "fast" speed. The extract was filtered through a 0.2 um filter into an autosampler tube. The filtered extracts were then used in HPLC analysis described below.

Leaf extracts were prepared by mixing 30-50 mg of leaf tissue with 1 g microbeads and freezing in liquid nitrogen until extraction. For extraction, 500 ul 1% pyrogallol in ethanol was added to the leaf/bead mixture and shaken for 1 minute on a Beadbeater (Biospec) on "fast" speed. The resulting mixture was centrifuged for 4 minutes at 14,000 rpm and filtered as described above prior to HPLC analysis.

HPLC was performed on a Zorbax silica HPLC column (4.6 mm X 250 mm) with a fluorescent detection, an excitation at 290 nm, an emission at 336 nm, and bandpass and slits. Solvent A was hexane and solvent B was methyl-t-butyl ether. The injection volume was 20 ul, the flow rate was 1.5 ml/min, the run time was 12 min (40°C) using the gradient (Table 5):

**Table 5:**

<u>Time</u>	<u>Solvent A</u>	<u>Solvent B</u>
0 min.	90%	10%
10 min.	90%	10%
11 min.	25%	75%
12 min.	90%	10%

Tocopherol standards in 1% pyrogallol/ ethanol were also run for comparison (alpha tocopherol, gamma tocopherol, beta tocopherol, delta tocopherol, and tocopherol (tocol) (all from Matreya).

Standard curves for alpha, beta, delta, and gamma tocopherol were calculated using Chemstation software. The absolute amount of component x is: Absolute amount of x=

$\text{Response}_x \times \text{RF}_x \times \text{dilution factor}$  where  $\text{Response}_x$  is the area of peak  $x$ ,  $\text{RF}_x$  is the response factor for component  $x$  ( $\text{Amount}_x/\text{Response}_x$ ) and the dilution factor is 500 ul. The ng/mg tissue is found by: total ng component/mg plant tissue.

Results of the HPLC analysis of seed extracts of transgenic *Arabidopsis* lines containing pMON10822 for the expression of ATAT2 from the napin promoter are provided in Figure 24.

HPLC analysis results of *Arabidopsis* seed tissue expressing the ATAT2 sequence from the napin promoter (pMON10822) demonstrates an increased level of tocopherols in the seed. Total tocopherol levels are increased as much as 50 to 60% over the total tocopherol levels of non-transformed (wild-type) *Arabidopsis* plants (Figure 24).

Furthermore, increases of particular tocopherols are also increased in transgenic *Arabidopsis* plants expressing the ATAT2 nucleic acid sequence from the napin promoter. Levels of delta tocopherol in these lines are increased greater than 3 fold over the delta tocopherol levels obtained from the seeds of wild type *Arabidopsis* lines. Levels of gamma tocopherol in transgenic *Arabidopsis* lines expressing the ATAT2 nucleic acid sequence are increased as much as about 60% over the levels obtained in the seeds of non-transgenic control lines. Furthermore, levels of alpha tocopherol are increased as much as 3 fold over those obtained from non-transgenic control lines.

Results of the HPLC analysis of seed extracts of transgenic *Arabidopsis* lines containing pMON10803 for the expression of ATAT2 from the enhanced 35S promoter are provided in Figure 25.

All publications and patent applications mentioned in this specification are indicative of the level of skill of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claim.

## Claims

What is Claimed is:

1. An isolated nucleic acid sequence encoding a prenyltransferase.
- 5        2. An isolated nucleic acid sequence according to Claim 1, wherein said prenyltransferase is selected from the group consisting of straight chain prenyltransferase and aromatic prenyltransferase.
3. An isolated DNA sequence according to Claim 1, wherein said nucleic acid sequence is isolated from a eukaryotic cell source.
4. An isolated DNA sequence according to Claim 3, wherein said eukaryotic cell source is  
10       selected from the group consisting of mammalian, nematode, fungal, and plant cells.
5. The DNA encoding sequence of Claim 4 wherein said prenyltransferase protein is from *Arabidopsis*.
6. The DNA encoding sequence of Claim 5 wherein said prenyltransferase protein is encoded by a sequence selected from the group consisting of the sequences of Figure 1.
- 15       7. The DNA encoding sequence of Claim 4 wherein said prenyltransferase protein is from corn.
8. The DNA encoding sequence of Claim 7 wherein said prenyltransferase protein is encoded by a sequence which includes the EST of the sequences of Figure 3.
9. The DNA encoding sequence of Claim 4 wherein said prenyltransferase protein is from  
20       soybean.
10. The DNA encoding sequence of Claim 9 wherein said prenyltransferase protein is encoded by a sequence which includes the ESTs of the group consisting of the sequences of Figure 2 and Figure 9.
11. An isolated DNA sequence according to Claim 1, wherein said nucleic acid sequence is  
25       isolated from a prokaryotic cell source.
12. An isolated DNA sequence according to Claim 11, wherein said prokaryotic source is *Synechocystis*.
13. A nucleic acid construct comprising as operably linked components, a transcriptional initiation region functional in a host cell, a nucleic acid sequence encoding prenyltransferase, and a  
30       transcriptional termination region.
14. A nucleic acid construct according to Claim 13, wherein said nucleic acid sequence encoding prenyltransferase is obtained from an organism selected from the group consisting of a eukaryotic organism and a prokaryotic organism.

15. A nucleic acid construct according to Claim 14, wherein said nucleic acid sequence encoding prenyltransferase is obtained from a plant source.

16. A nucleic acid construct according to Claim 15, wherein said nucleic acid sequence encoding prenyltransferase is obtained from a source selected from the group consisting of

5 *Arabidopsis*, soybean and corn.

17. A nucleic acid construct according to Claim 13, wherein said nucleic acid sequence encoding prenyltransferase is obtained from *Synechocystis*.

18. A plant cell comprising the construct of Claim 13.

19. A method for the alteration of the tocopherol content in a host cell, comprising;  
10 transforming said host cell with a construct comprising as operably linked components, a transcriptional initiation region functional in a host cell, a nucleic acid sequence encoding prenyltransferase, and a transcriptional termination region.

20. The method according to Claim 19, wherein said host cell is selected from the group consisting of a prokaryotic cell and a eukaryotic cell.

15 21. The method according to Claim 20, wherein said prokaryotic cell is *Synechocystis*.

22. The method according to Claim 20, wherein said eukaryotic cell is a plant cell.

23. The method according to Claim 22, wherein said plant cell is obtained from a plant selected from the group consisting of *Arabidopsis*, soybean, and corn.

24. A method for producing a tocopherol compound of interest in a host cell, said method  
20 comprising obtaining a transformed host cell, said host cell having and expressing in its genome:  
a construct having a DNA sequence encoding a prenyltransferase operably linked to a transcriptional initiation region functional in a host cell,  
wherein said prenyltransferase is involved in the synthesis of tocopherols.

25 25. The method according to Claim 24, wherein said host cell is selected from the group consisting of a prokaryotic cell and a eukaryotic cell.

26. The method according to Claim 25, wherein said prokaryotic cell is *Synechocystis*.

27. The method according to Claim 24, wherein said eukaryotic cell is a plant cell.

28. The method according to Claim 27, wherein said plant cell is obtained from a plant selected from the group consisting of *Arabidopsis*, soybean, and corn.

30 29. A method for increasing the biosynthetic flux in cell from a host cell toward tocopherol production, said method comprising transforming said host cell with a construct comprising as operably linked components, a transcriptional initiation region functional in a

host cell, a DNA encoding a prenyltransferase involved in the synthesis of tocopherols, and a transcriptional termination region.

30. The method according to Claim 29, wherein said host cell is selected from the group consisting of a prokaryotic cell and a eukaryotic cell.

5 31. The method according to Claim 30, wherein said prokaryotic cell is *Synechocystis*.

32. The method according to Claim 30, wherein said eukaryotic cell is a plant cell.

33. The method according to Claim 32, wherein said plant cell is obtained from a plant selected from the group consisting of *Arabidopsis*, soybean, and corn.

10




Figure 1






Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8






Figure 9




Figure 10




Figure 11




Figure 12




Figure 13




Figure 14




Figure 15






Figure 16




Figure 17




Figure 18




Figure 19




Figure 20

SLR1736 : MATIQAFWR---FSRPHTIIGTTLS\* 20 40 60 80  
 SLR0926 : MVAQHPSS---PPLWLTIIYLRWHPK\* 60 80  
 SLL1899 : MVTSMKIHRQDSMGVCKSYQLTKE\* 60 80  
 SLR0056 : MSDAONTGQ--NQAKRQLLGMKGAAPGESSIWKIRLQMKPITWIPLIHWGVCCGAASSGGYIWSVEDFELKALTCMLISGPIMTGVTQT\* 60 80  
 SLR1518 : MTESSPLAP--STAPATRKLWAAIKP\* 60 80  
 M 100 120 140 160 180  
 SLR1736 : NQIWDNDIERINKPNLPANGDFSIAQGRWIVGICGVASIAIAWGLGLWGLTVGIS--LITGTAYSVPFRLKRFSLLAALCITTVRGI\* 100 120 140 160 180  
 SLR0926 : NDLWDRDIDPOVERTKORSLARAL-SVQVGICVALVA-LICCAACAFYITPLS-----FWLCVAAAPVVAAYPGARRVFPVPOAVL-SL\* 100 120 140 160 180  
 SLL1899 : NCLYDQDITIEYMLRTKARIPAGKV-OPRHAIIFALALEVSEFALLATFVNLVSGC--LALSEIVFYMLVYTHWLKRHTAQNIVIG-GA\* 100 120 140 160 180  
 SLR0056 : NLFYDRDIDAINEPYRPESGAIISVPOVVTQHLILLVAGIGVAYGLDVMAQHDFFIMMVITLGGAFAYAYSAPPLKLKQNGWLGNY-AL\* 100 120 140 160 180  
 SLR1518 : NDVEDSDITGIDVRKAHSMVNLGTGRNLYFLISNFFELFAGVGLMSMSWRAQDWT-----VLEELIGVAFLFGYTYQGPPFRRLGYLGLGELIC\* 100 120 140 160 180  
 N 5D Did 1 a 6 6 1 6  
 SLR1736 : VVNLGLFDFERIGLGYPPPTLITPIANLILFILPPTVAIAIFKDMPDMEGDRQFKIQTLTIQIGKONVFRGTLGLLTG-CYIAMAIIWGLWA\* 200 220 240 260  
 SLR0926 : AWGFAVITSMS---AVRGDIDATVNLGATVFWNLGFDTVYAMADREDDRRIGVNSALFFG-CYVGEAVGIEFFA--LITIGCLFYUGMI\* 200 220 240 260  
 SLL1899 : AGSIPPTVGMIA---AVTGDLSWTPWNLFALITFWPPHFWALAIMIKDDYAGVNVPMHPVIAEGEKTVSQIWIYYSL--DVVPPSLLVYP\* 200 220 240 260  
 SLR0056 : GASVIAIPWMAAG-HALFCTLNPTAMLLNIIYSIAGEGHVVNDFKSVGEGDRQGLGKSLPVMFG-IGTAAWLCVYMI--DVFQAGIAGYLI\* 200 220 240 260  
 SLR1518 : LITEGPIAIAAAYYSQSQSFWNLTPSVFVGISTAIILFCSHFHQVEDDLAAGKKSPIVRLG-TKLGSQVLTLSVVSLYITITALIGVLCH\* 200 220 240 260  
 1 vl 1 vl t e d 6 G  
 SLR1736 : AMPLNTATITVSHDCCLALLMWRSRDVHLESKTEHASFYQIWKDFLEYLYPLALWLPNFSNTIF-----: 308  
 SLR0926 : LMLNPHYMSIAATAVGWVIOYIQLSAPTEPKLYGQ--IIGQNVIIIGFVLLAGMLLGW-----: 292  
 SLL1899 : LHQLGGLLYAATATIGGOFILVKAWQLKQAPGDRDLARG-LFKFSBYLMLLCLAMVLDSPVTHQLVAQMGTLLG: 316  
 SLR0056 : YVHQQLYATITVLLLTIPQITFQDMYFLRNPLENDVKYQ-ASAQPFLVFGMLATSEALGHAGI-----: 324  
 SLR1518 : QAPWQTLLLTIASLPWAVQVIRHVGGYQYHDQPEQVSNCKFIAVNLHFFSGMIMAAAGYGWAGIG-----: 307  
 280 \* 300 320 340 p 6

Figure 21

	*	20	*	40	*	60	*	80	
ATPT2	----	MESLLSSSLVSAAGGFCWKKNLKLHSLSEIRVLRCDSSKVVAKPKFRNNLVRPDPGQSSLLLYPKHKSRFRVNATAGQ	----	----	----	----	----	----	80
SLR1736	----	-----	-----	-----	-----	-----	-----	-----	-
ATPT3	----	MAFFGLSRVSRLLKSSVSVTPSSSALLQSQHKSLSNPVTTHYTNPFCKYPSWNDNYQVWSKGRHLHQEKFFGVGWNRYRLICGMSSS	----	----	----	----	----	----	89
SLR0926	----	-----	-----	-----	-----	-----	-----	-----	-
ATPT4	----	MWRRSVVYRFSRISVSSSLPNRPLIPWSRELCAVNSFSQP	----	----	----	----	----	----	67
SLR1899	----	-----	-----	-----	-----	-----	-----	-----	4
ATPT12	----	-----	-----	-----	-----	-----	-----	-----	63
SLR0056	----	-----	-----	-----	-----	-----	-----	-----	4
ATPT8	----	-----	-----	-----	-----	-----	-----	-----	4
SLR1518	----	-----	-----	-----	-----	-----	-----	-----	4

	*	100	*	120	*	140	*	160	*	1
ATPT2	----	PEAFDSNSKQK	----	SFRDSIDAFYR	----	FSRPHTVIGTVLSLS	----	VSFLAVEKVS	----	DISPLFTGILE
SLR1736	----	-----	-----	MATIQAFWR	----	FSRPHTVIGTVLSLS	----	LLTILGDGN	----	SVNSPASUDLVFG
ATPT3	----	SSVLEGPCKDDKEKSDGVVKKASW	----	PLWITIIYL	----	LRWHKPAGRILMIPALMA	----	IALAADPGS	----	LPSFKYMALFGC
SLR0926	----	-----	-----	-----	-----	-----	-----	VCLAAQ	----	G-LPPLPLUGTIAL
ATPT4	----	TAAATATATTG	----	EISSRVAALAGLGHYAR	----	CYWELSKAKISMLVATSG	----	TGYILGTGNAAISFPCLCYTCAG	----	138
SLR1899	----	TKIHRQHDMSG	----	AVCKSYQQLTKP	----	RIIPLLLTTAASMI	----	ASEGR	----	VDPKHLITLLG
ATPT12	----	DKVKSQTPDKAP	----	AGGSSINQLLTKGAS	----	QETNKWKIRLQTKPVTWPLVWGVVCC	----	AAASGNFHWTPEDVAKSLC	----	139
SLR0056	----	QNT-GQNQAKA	----	ROLLGKGAAP	----	GESSIWKIRLQMKPITWPLIWGVVCC	----	AAASGGYIWSVEDFKALTC	----	73
ATPT8	----	EVPKLASAAEY	----	FFKRGVQKQF	----	RSTILLMATAALNRVP	----	EALIGEST	----	DIVTSELVRQR
SLR1518	----	SPLAPSTAPAT	----	RKLWAAAIKP	----	PMYTMVAVVPITVG	----	SAVAYGLTG	----	QWHDVFTIFLL

	*	200	*	220	*	240	*	260		
ATPT2	----	AVVAALMNIYIVGLNQIS	----	VEIKVKNKPYLPLASGEYS	----	NTGIAHVASFS	----	MSFWLGIWVGSWPLFWALFVSF	----	MLGTAYS-INPILR
SLR1736	----	AWLACLLGNVYIVGLNQIS	----	MDVILIRINKPNLPLANGDFS	----	QAQGRWGLCGVAS	----	AIATWGLG-LWLGLTVGLSL	----	IGTAY---SMPVVR
ATPT3	----	GAILL---	----	ERGAGCTINDELLO	----	QIDTKVDRTKLRPIASGLIT	----	PFQCGFLGLQLLGLG	----	ILLQLNNYSRVLGAS---
SLR0926	----	GTICA---	----	TSGLGCVVNDLMDRDL	----	POVERTKQRPAAARALS	----	VQVGLGVALVALCAG	----	LFYLTPLSFMLCVH---
ATPT4	----	TMMI---	----	AASANSINQIHEISND	----	SKMKRTMLRPLPSGRISVPHAV	----	AWATAGASGACL	----	LASKTNMIAAGLASAN---
SLR1899	----	GTIA---	----	AASAQTLNCLYTOIIL	----	VEMLRTRARIPAGKVOPRHA	----	FALAGVLSFNL	----	LATFVNVSGLALSG---
ATPT12	----	MMMSGPCITGYTOTINDW	----	YDRIDAINEPYRPIESGAI	----	SEPEVITQWVLGLGG	----	GIAGILD-VWAGHTTPTVFYALG	----	---GSLISY
SLR0056	----	MLISGPLMTGYTQTLNDF	----	YDRIDAINEPYRPIESGAI	----	SVQVVTQILILVAG	----	GVAYGLD-VWAQHDFFMM	----	MLTLG---GAFWAY
ATPT8	----	GIAE---	----	TEMIHVASLHDDVLEDD	----	ADADTRRGVGSNNVMGNKMSV	----	ADGDFESRACGAL	----	AAALKNTEVVALATAVEHLVTGETM
SLR1518	----	SATA---	----	TIWINLSNDVFDSDT	----	GIDVRKAHSVUNLTGNNRNLVFL	----	SNFFLAGVGLGLMSMS	----	WRAQDWTVLELIGVA---

Figure 22 1/2



```

ATPT2 : WKR-FALVAAMCILA VRAII QIAFYH-IQTHVFGRPITRPIEFATAFMSF SVVIALFKDIPDI GKIK-FGRSFSVTLG-QKR : 313
SLR1736 : LKR-FSLLAALCILT VRGIVNLGLFIF-FRIGLGYPTTITPMMV-TLFLVFAVAIAAFKDVDPDMGRQ-FKQIQTTLQIG-KQN : 218
ATPT3 : SVP-LMKRFTFWPQAFGLLTENWGALHG--WT--ATK2SIAPSIVLPYLSGVC--LVYDIIYAHQDKDDVK--GVKSTALRFG-DNT : 328
SLR0926 : AVP-GAKRVFPVPQLVLSIANGFAVETIS--WS--AVTEDLDATW--WGATVF--LGFDIVYAMADRDDR--GVNNSALFFC-QYV : 213
ATPT4 : VPT-PLKQLHPINTWVGAVVGAIPPEIG--WA--AASQISYNSMI--PPAALYFQIPIPHMALAHLCRNDYAA-GGYKMSIFDP--S : 294
SLL1899 : VTHWLKRHTAQNIIVIGGAAGSIPPEIG--WA--AVTEDLDATW--FALIFLWTPPHFWALALMIKDDYAQ--VNVPMIPVIAEET : 220
ATPT12 : IYS-APPLKQKONGWVGNFAGASYISLPWAGQALFETIPDQV--TLLYSIAGIGIADVNDFKSVEGDRA--LGIQSIIPVAFG-TET : 308
SLR0056 : IYS-APPLKQKONGWVGNFAGASYISLPWAGHALFETINPTW--TLLIYSLAGIGIADVNDFKSVEGDRA--LGIQSIIPVAFG-TET : 242
ATPT8 : EITSSTEQRYSMDYYMQTTYKKTASISNSCKAVAVLTGQAEVAVLAFEYGRNLGLAFQIIDDILDFTGTSASLKGSSDIRH--GV : 231
SLR1518 : TYQGPFRRLGGLGELICLITFGPFAI-AAAYYSQSQSFSWNHET-PSVFGVIGISTAIILFCSHFHQVEDDLA-AKKKSPIVRLG-TKL : 223

```

```

ATPT2 : VFWTCVTLQOMAYAVAIIVGATSPFIWSKVISVGHVITATTIWARAKSVLSSKTEITS---CYMFIWKIFUYAEYGLLPFLK----- : 393
SLR1736 : VFRGTLILITGTCYLAMATWGLWAAMPLNTAFIIVSHLC--LABLWRSRDVHLESKTEIAS---FYQIWKIFFLEYGLYPIALWLPNFS : 304
ATPT3 : KLWITGFGTASIGFIALSGFSADLGWQYASIAASGQIGWQIGTADLSSGADCS---RKIVSNKWFGAIFFGVGRSFQ- : 407
SLR0926 : GEAVGTFEFTIGCFYIGMLMLNPLYWLSJAAAI--GVWIIQYIQLSAPTPEP-KIY---GOIFGQNIIGFVHAGMLLGWL--- : 292
ATPT4 : GKRFAAVALRNCFYIPIGFAIDWGLTSSWFCUESTLTALAATAAFSFRDRTMHKA---RKMFFHASULFLPVMFSGILLHRVSND : 379
SLL1899 : VSQIWIYYSLLVVPFSLHVVPLHQLGILYLATAIIT--GGQFLVKAWQLKQAPGDRDIA---RGLKFSFYLMIECLAMVDSLPT : 303
ATPT12 : AKWICVGAIDITQLSVAGYLASGKPYALALVALI--IPQVVFQFKYFLKDPKYDYK---YQASAPFLVLGIFVTALASQH--- : 387
SLR0056 : AAWICVIMIDVFOAGIAGYLYVHQQLYATIVLLI--IPQITFQDMYFLRNPLENDVK---YQASAPFLVFGMLATGIALGHAGI- : 324
ATPT8 : ITAPILFAMEEFPQREIVDOVEKDPNVDIAIEYIGKSGKQORARELANEHLAAAGSLPETDNEDVKRSRRRAIDITHRVITRN : 320
SLR1518 : GSQVLTLSVSLYLITATAGVLCCHQAPWQTLTLLTASLPWAVQLEHRVGOYHDOPEQVSNCK---FIAVNLHFFSGMIMAAGYGWAGLG-- : 307


```

```

ATPT2 : ----- : -
SLR1736 : NTIF----- : 308
ATPT3 : ----- : -
SLR0926 : ----- : -
ATPT4 : NQQOLVEEAGLTNSVSGEVKQRRKRVAQPPVAVASAAFPFLPAPSFYSP : 431
SLL1899 : --HQLVAQMGTLLG----- : 316
ATPT12 : ----- : -
SLR0056 : ----- : -
ATPT8 : K----- : 321
SLR1518 : ----- : -

```

Figure 22 2/2



*Synechocystis* 6803 wild type      *Synechocystis* slr1736 knockout

Figure 23




Figure 24




Figure 25

## SEQUENCE LISTING

5 <110> Calgene LLC  
 <120> Nucleic Acid Sequences Involved in  
 Tocopherol Synthesis  
 <130> 17133/00/WO  
 10 <150> 60/129,899  
 <151> 1999-04-15  
 <150> 60/146,461  
 15 <151> 1999-07-30  
 <160> 94  
 <170> FastSEQ for Windows Version 4.0  
 20 <210> 1  
 <211> 1182  
 <212> DNA  
 <213> Arabidopsis sp  
 25 <400> 1  
 atggaggtctc tgctctctag ttctttctctt gtttccgctg ctgggtgggtt ttgttggaag 60  
 aagcagaatc taaagctcca ctctttatca gaaatccgag ttctgcgttg tgattcgagt 120  
 aaagttgtcg caaaaccgaa gtttaggaac aatcttggtta ggcctgatgg tcaaggatct 180  
 30 tcattggttg tgtatccaaa acataagtcg agatttcggg ttaatgccac tgcgggtcag 240  
 cctgaggctt tcgactcgaa tagcaaacag aagtctttta gagactcgtt agatgcgttt 300  
 tacagggttt ctaggcctca tacagttatt ggcacagtgc ttagcatttt atctgtatct 360  
 ttcttagcag tagagaaggt ttctgatata tctcctttac ttttcaactgg catcttggag 420  
 gctgttggtg cagctctcat gatgaacatt tacatagttg ggctaaatca gttgtctgat 480  
 35 gttgaaatag ataagggttaa caagccctat ctccatttg catcaggaga atattctggt 540  
 aacaccggca ttgcaatagt agcttccttc tccatcatga gtttctggct tgggtggatt 600  
 gttgggtcat ggccattgtt ctgggctctt tttgtgagtt tcatgctcgg tactgcatac 660  
 tctatcaatt tgccactttt acggttgaaa agatttgcac tggttgcagc aatgtgtatc 720  
 ctgcgtgtcc gagctattat tgttcaaate gcctttttatc tacatattca gacacatgtg 780  
 40 tttggaagac caatcttggt cactaggcct ctatttttcg ccaactgcgtt tatgagcttt 840  
 ttctctgtcg ttattgcatt gtttaaggat atacctgata tcgaagggga taagatattc 900  
 ggaatccgat cattctctgt aactctgggt cagaaacggg tgttttggac atgtgttaca 960  
 ctacttcaaa tggcttaacg tgttgcaatt ctagtggag ccacatctcc attcatatgg 1020  
 agcaaagtca tctcgggttg gggtcattgt atactcgcaa caactttgtg ggctcgagct 1080  
 45 aagtccgttg atctgagtag caaaaccgaa ataacttcat gttatatgtt catatggaag 1140  
 ctcttttatg cagagtactt gctgttacct tttttgaagt ga 1182

&lt;210&gt; 2

&lt;211&gt; 393

&lt;212&gt; PRT

5 &lt;213&gt; Arabidopsis sp

&lt;400&gt; 2

	Met	Glu	Ser	Leu	Leu	Ser	Ser	Ser	Ser	Leu	Val	Ser	Ala	Ala	Gly	Gly
	1				5					10					15	
10	Phe	Cys	Trp	Lys	Lys	Gln	Asn	Leu	Lys	Leu	His	Ser	Leu	Ser	Glu	Ile
				20					25					30		
	Arg	Val	Leu	Arg	Cys	Asp	Ser	Ser	Lys	Val	Val	Ala	Lys	Pro	Lys	Phe
				35				40					45			
	Arg	Asn	Asn	Leu	Val	Arg	Pro	Asp	Gly	Gln	Gly	Ser	Ser	Leu	Leu	Leu
15		50					55					60				
	Tyr	Pro	Lys	His	Lys	Ser	Arg	Phe	Arg	Val	Asn	Ala	Thr	Ala	Gly	Gln
	65					70				75					80	
	Pro	Glu	Ala	Phe	Asp	Ser	Asn	Ser	Lys	Gln	Lys	Ser	Phe	Arg	Asp	Ser
					85					90				95		
20	Leu	Asp	Ala	Phe	Tyr	Arg	Phe	Ser	Arg	Pro	His	Thr	Val	Ile	Gly	Thr
				100					105					110		
	Val	Leu	Ser	Ile	Leu	Ser	Val	Ser	Phe	Leu	Ala	Val	Glu	Lys	Val	Ser
				115				120					125			
	Asp	Ile	Ser	Pro	Leu	Leu	Phe	Thr	Gly	Ile	Leu	Glu	Ala	Val	Val	Ala
25		130					135					140				
	Ala	Leu	Met	Met	Asn	Ile	Tyr	Ile	Val	Gly	Leu	Asn	Gln	Leu	Ser	Asp
	145					150					155				160	
	Val	Glu	Ile	Asp	Lys	Val	Asn	Lys	Pro	Tyr	Leu	Pro	Leu	Ala	Ser	Gly
					165					170					175	
30	Glu	Tyr	Ser	Val	Asn	Thr	Gly	Ile	Ala	Ile	Val	Ala	Ser	Phe	Ser	Ile
				180					185					190		
	Met	Ser	Phe	Trp	Leu	Gly	Trp	Ile	Val	Gly	Ser	Trp	Pro	Leu	Phe	Trp
				195				200					205			
	Ala	Leu	Phe	Val	Ser	Phe	Met	Leu	Gly	Thr	Ala	Tyr	Ser	Ile	Asn	Leu
35		210					215					220				
	Pro	Leu	Leu	Arg	Trp	Lys	Arg	Phe	Ala	Leu	Val	Ala	Ala	Met	Cys	Ile
	225					230					235				240	
	Leu	Ala	Val	Arg	Ala	Ile	Ile	Val	Gln	Ile	Ala	Phe	Tyr	Leu	His	Ile
					245					250					255	
40	Gln	Thr	His	Val	Phe	Gly	Arg	Pro	Ile	Leu	Phe	Thr	Arg	Pro	Leu	Ile
				260					265					270		
	Phe	Ala	Thr	Ala	Phe	Met	Ser	Phe	Phe	Ser	Val	Val	Ile	Ala	Leu	Phe
				275				280					285			
	Lys	Asp	Ile	Pro	Asp	Ile	Glu	Gly	Asp	Lys	Ile	Phe	Gly	Ile	Arg	Ser
45		290					295					300				
	Phe	Ser	Val	Thr	Leu	Gly	Gln	Lys	Arg	Val	Phe	Trp	Thr	Cys	Val	Thr

3

	Met	Ala	Phe	Phe	Gly	Leu	Ser	Arg	Val	Ser	Arg	Arg	Leu	Leu	Lys	Ser	
	1				5					10					15		
	Ser	Val	Ser	Val	Thr	Pro	Ser	Ser	Ser	Ser	Ala	Leu	Leu	Gln	Ser	Gln	
				20					25					30			
5	His	Lys	Ser	Leu	Ser	Asn	Pro	Val	Thr	Thr	His	Tyr	Thr	Asn	Pro	Phe	
			35					40					45				
	Thr	Lys	Cys	Tyr	Pro	Ser	Trp	Asn	Asp	Asn	Tyr	Gln	Val	Trp	Ser	Lys	
		50					55					60					
	Gly	Arg	Glu	Leu	His	Gln	Glu	Lys	Phe	Phe	Gly	Val	Gly	Trp	Asn	Tyr	
10	65					70					75					80	
	Arg	Leu	Ile	Cys	Gly	Met	Ser	Ser	Ser	Ser	Ser	Val	Leu	Glu	Gly	Lys	
				85						90					95		
	Pro	Lys	Lys	Asp	Asp	Lys	Glu	Lys	Ser	Asp	Gly	Val	Val	Val	Lys	Lys	
				100					105					110			
15	Ala	Ser	Trp	Ile	Asp	Leu	Tyr	Leu	Pro	Glu	Glu	Val	Arg	Gly	Tyr	Ala	
			115					120					125				
	Lys	Leu	Ala	Arg	Leu	Asp	Lys	Pro	Ile	Gly	Thr	Trp	Leu	Leu	Ala	Trp	
		130					135					140					
	Pro	Cys	Met	Trp	Ser	Ile	Ala	Leu	Ala	Ala	Asp	Pro	Gly	Ser	Leu	Pro	
20	145					150					155					160	
	Ser	Phe	Lys	Tyr	Met	Ala	Leu	Phe	Gly	Cys	Gly	Ala	Leu	Leu	Leu	Arg	
				165						170					175		
	Gly	Ala	Gly	Cys	Thr	Ile	Asn	Asp	Leu	Leu	Asp	Gln	Asp	Ile	Asp	Thr	
				180					185					190			
25	Lys	Val	Asp	Arg	Thr	Lys	Leu	Arg	Pro	Ile	Ala	Ser	Gly	Leu	Leu	Thr	
			195					200					205				
	Pro	Phe	Gln	Gly	Ile	Gly	Phe	Leu	Gly	Leu	Gln	Leu	Leu	Leu	Gly	Leu	
		210					215					220					
	Gly	Ile	Leu	Leu	Gln	Leu	Asn	Asn	Tyr	Ser	Arg	Val	Leu	Gly	Ala	Ser	
30	225					230					235					240	
	Ser	Leu	Leu	Leu	Val	Phe	Ser	Tyr	Pro	Leu	Met	Lys	Arg	Phe	Thr	Phe	
				245						250				255			
	Trp	Pro	Gln	Ala	Phe	Leu	Gly	Leu	Thr	Ile	Asn	Trp	Gly	Ala	Leu	Leu	
				260					265					270			
35	Gly	Trp	Thr	Ala	Val	Lys	Gly	Ser	Ile	Ala	Pro	Ser	Ile	Val	Leu	Pro	
			275					280					285				
	Leu	Tyr	Leu	Ser	Gly	Val	Cys	Trp	Thr	Leu	Val	Tyr	Asp	Thr	Ile	Tyr	
		290					295					300					
	Ala	His	Gln	Asp	Lys	Glu	Asp	Asp	Val	Lys	Val	Gly	Val	Lys	Ser	Thr	
40	305					310					315					320	
	Ala	Leu	Arg	Phe	Gly	Asp	Asn	Thr	Lys	Leu	Trp	Leu	Thr	Gly	Phe	Gly	
				325						330				335			
	Thr	Ala	Ser	Ile	Gly	Phe	Leu	Ala	Leu	Ser	Gly	Phe	Ser	Ala	Asp	Leu	
				340					345					350			
45	Gly	Trp	Gln	Tyr	Tyr	Ala	Ser	Leu	Ala	Ala	Ala	Ser	Gly	Gln	Leu	Gly	
			355					360					365				



Trp Gln Ile Gly Thr Ala Asp Leu Ser Ser Gly Ala Asp Cys Ser Arg  
 370 375 380  
 Lys Phe Val Ser Asn Lys Trp Phe Gly Ala Ile Ile Phe Ser Gly Val  
 385 390 395 400  
 5 Val Leu Gly Arg Ser Phe Gln  
 405

<210> 5

<211> 1296

10 <212> DNA

<213> Arabidopsis sp

<400> 5

	atgtggcgaa gatctgttgt ttctcgttta tcttcaagaa tctctgtttc ttcttcgtta	60
15	ccaaacccta gactgattcc ttgggtccgc gaattatgtg ccgttaatag cttctcccag	120
	cctccggtct cgacggaatc aactgctaag ttagggatca ctggtgtag atctgatgcc	180
	aatcgagttt ttgccactgc tactgccgcc gctacagcta cagctaccac cggtgagatt	240
	tcgtctagag ttgcggcttt ggctggatta gggcatcact acgctcgttg ttattgggag	300
	ctttctaaag ctaaacttag tatgcttgtg gttgcaactt ctggaactgg gtatattctg	360
20	ggtaggggaa atgctgcaat tagcttcccg gggctttgtt acacatgtgc aggaaccatg	420
	atgattgctg catctgctaa ttccttgaat cagatttttg agataagcaa tgattctaa	480
	atgaaaagaa cgatgctaag gccattgcct tcaggacgta ttagtggtcc acacgctgtt	540
	gcatgggcta ctattgctgg tgcttctggg gcttgtttgt tggccagcaa gactaatatg	600
	ttggctgctg gacttgcac tgccaatctt gtactttatg cgtttgttta tactccgttg	660
25	aagcaacttc accctatcaa tacatgggtt ggcgctgttg ttggtgctat cccacccttg	720
	cttgggtggg cggcagcgtc tggtcagatt tcatacaatt cgatgattct tccagctgct	780
	ctttactttt ggcagatacc tcattttatg gcccttgac atctctgccg caatgattat	840
	gcagctggag gttacaagat gttgtcactc tttgatccgt caggggaagag aatagcagca	900
	gtggctctaa ggaactgctt ttacatgatc cctctcggtt tcatcgcta tgactggggg	960
30	ttaacctcaa gttgggttttg cctcgaatca acacttctca cactagcaat cgctgcaaca	1020
	gcattttcat tctaccgaga ccggaccatg cataaagcaa ggaaaatgtt ccatgccagt	1080
	cttctcttcc ttctgtttt catgtctggg cttcttctac accgtgtctc taatgataat	1140
	cagcaacaac tcgtagaaga agccggatta acaaattctg tatctggtga agtcaaaact	1200
	cagaggcgaa agaaacgtgt ggctcaacct ccggtggctt atgcctctgc tgcaccgttt	1260
35	cctttctctc cagctccttc cttctactct ccatga	1296

<210> 6

<211> 431

<212> PRT

40 <213> Arabidopsis sp

<400> 6

	Met Trp Arg Arg Ser Val Val Tyr Arg Phe Ser Ser Arg Ile Ser Val	
	1 5 10 15	
45	Ser Ser Ser Leu Pro Asn Pro Arg Leu Ile Pro Trp Ser Arg Glu Leu	
	20 25 30	

	Cys	Ala	Val	Asn	Ser	Phe	Ser	Gln	Pro	Pro	Val	Ser	Thr	Glu	Ser	Thr	
			35					40					45				
	Ala	Lys	Leu	Gly	Ile	Thr	Gly	Val	Arg	Ser	Asp	Ala	Asn	Arg	Val	Phe	
		50					55				60						
5	Ala	Thr	Ala	Thr	Ala	Ala	Ala	Thr	Ala	Thr	Ala	Thr	Thr	Gly	Glu	Ile	
	65					70					75				80		
	Ser	Ser	Arg	Val	Ala	Ala	Leu	Ala	Gly	Leu	Gly	His	His	Tyr	Ala	Arg	
					85				90					95			
10	Cys	Tyr	Trp	Glu	Leu	Ser	Lys	Ala	Lys	Leu	Ser	Met	Leu	Val	Val	Ala	
				100					105					110			
	Thr	Ser	Gly	Thr	Gly	Tyr	Ile	Leu	Gly	Thr	Gly	Asn	Ala	Ala	Ile	Ser	
			115					120					125				
	Phe	Pro	Gly	Leu	Cys	Tyr	Thr	Cys	Ala	Gly	Thr	Met	Met	Ile	Ala	Ala	
		130					135					140					
15	Ser	Ala	Asn	Ser	Leu	Asn	Gln	Ile	Phe	Glu	Ile	Ser	Asn	Asp	Ser	Lys	
	145					150					155				160		
	Met	Lys	Arg	Thr	Met	Leu	Arg	Pro	Leu	Pro	Ser	Gly	Arg	Ile	Ser	Val	
					165				170					175			
20	Pro	His	Ala	Val	Ala	Trp	Ala	Thr	Ile	Ala	Gly	Ala	Ser	Gly	Ala	Cys	
				180				185						190			
	Leu	Leu	Ala	Ser	Lys	Thr	Asn	Met	Leu	Ala	Ala	Gly	Leu	Ala	Ser	Ala	
			195					200					205				
	Asn	Leu	Val	Leu	Tyr	Ala	Phe	Val	Tyr	Thr	Pro	Leu	Lys	Gln	Leu	His	
		210					215					220					
25	Pro	Ile	Asn	Thr	Trp	Val	Gly	Ala	Val	Val	Gly	Ala	Ile	Pro	Pro	Leu	
	225					230					235				240		
	Leu	Gly	Trp	Ala	Ala	Ala	Ser	Gly	Gln	Ile	Ser	Tyr	Asn	Ser	Met	Ile	
				245					250					255			
30	Leu	Pro	Ala	Ala	Leu	Tyr	Phe	Trp	Gln	Ile	Pro	His	Phe	Met	Ala	Leu	
				260				265					270				
	Ala	His	Leu	Cys	Arg	Asn	Asp	Tyr	Ala	Ala	Gly	Gly	Tyr	Lys	Met	Leu	
			275				280					285					
	Ser	Leu	Phe	Asp	Pro	Ser	Gly	Lys	Arg	Ile	Ala	Ala	Val	Ala	Leu	Arg	
		290					295					300					
35	Asn	Cys	Phe	Tyr	Met	Ile	Pro	Leu	Gly	Phe	Ile	Ala	Tyr	Asp	Trp	Gly	
	305					310					315				320		
	Leu	Thr	Ser	Ser	Trp	Phe	Cys	Leu	Glu	Ser	Thr	Leu	Leu	Thr	Leu	Ala	
				325					330					335			
40	Ile	Ala	Ala	Thr	Ala	Phe	Ser	Phe	Tyr	Arg	Asp	Arg	Thr	Met	His	Lys	
				340				345						350			
	Ala	Arg	Lys	Met	Phe	His	Ala	Ser	Leu	Leu	Phe	Leu	Pro	Val	Phe	Met	
			355					360					365				
	Ser	Gly	Leu	Leu	Leu	His	Arg	Val	Ser	Asn	Asp	Asn	Gln	Gln	Gln	Leu	
		370				375						380					
45	Val	Glu	Glu	Ala	Gly	Leu	Thr	Asn	Ser	Val	Ser	Gly	Glu	Val	Lys	Thr	
	385					390					395				400		

Gln Arg Arg Lys Lys Arg Val Ala Gln Pro Pro Val Ala Tyr Ala Ser  
 405 410 415  
 Ala Ala Pro Phe Pro Phe Leu Pro Ala Pro Ser Phe Tyr Ser Pro  
 420 425 430

5

<210> 7  
 <211> 479  
 <212> DNA  
 <213> Arabidopsis sp

10

<400> 7  
 ggaaactccc ggagcacctg tttgcaggta ccgctaacct taatcgataa tttattttctc 60  
 ttgtcaggaa ttatgtaagt ctggtggaag gctcgcatac cattttttgca ttgccttttcg 120  
 ctatgatcgg gtttactttg ggtgtgatga gaccaggcgt ggcttttatgg tatggcgaaa 180  
 15 acccattttt atccaatgct gcattccctc ccgatgattc gttctttcat tcctatacag 240  
 gtatcatgct gataaaactg ttactgggtac tggttttgtat ggtatcagca agaagcgcg 300  
 cgatggcggt taaccgggtat ctcgacaggc attttgacgc gaagaaccgc cgtactgcca 360  
 tccgtgaaat acctgcgggc gtcataatctg ccaacagtgc gctgggtgttt acgataggct 420  
 gctgcgtggg attctgggtg gcctgttatt tcattaacac gatctgtttt tacctggcg 479

20

<210> 8  
 <211> 551  
 <212> DNA  
 <213> Arabidopsis sp

25

<220>  
 <221> misc\_feature  
 <222> (1)... (551)  
 <223> n = A,T,C or G

30

<400> 8  
 ttgtggctta caccttaatg agcatacgcc agnccattac ggctcgttaa tcggcgccat 60  
 ngccggngct gntgcaccgg tagtgggcta ctgcgcctg accaatcagc ttgatctagc 120  
 ggctcttatt ctgttttttaa ttttactgtt ctggcaaata ccgcattttt acgcgatttc 180  
 35 cattttcagg ctaaaagact tttcagcggc ctgtattccg gtgctgcca tcattaaaga 240  
 cctgcgctat accaaaatca gcatgctggg ttacgtgggc ttatttacac tggctgctat 300  
 catgccggcc ctcttagggg atgccgggtg gatttatggg atagcggcct taattttagg 360  
 cttgtattgg ctttatattg ccatacaagg attcaagacc gccgatgatc aaaaatggtc 420  
 tcgtaagatg tttggatctt cgatttttaat cattaccctc ttgtcggtaa tgatgcttgt 480  
 40 ttaaacttac tgcctcctga agtttatata tcgataattt cagcttaagg aggcttagtg 540  
 gttaattcaa t 551

<210> 9  
 <211> 297  
 45 <212> PRT  
 <213> Arabidopsis sp

<400> 9

	Met	Val	Leu	Ala	Glu	Val	Pro	Lys	Leu	Ala	Ser	Ala	Ala	Glu	Tyr	Phe
	1				5					10					15	
5	Phe	Lys	Arg	Gly	Val	Gln	Gly	Lys	Gln	Phe	Arg	Ser	Thr	Ile	Leu	Leu
				20					25					30		
	Leu	Met	Ala	Thr	Ala	Leu	Asn	Val	Arg	Val	Pro	Glu	Ala	Leu	Ile	Gly
			35					40					45			
	Glu	Ser	Thr	Asp	Ile	Val	Thr	Ser	Glu	Leu	Arg	Val	Arg	Gln	Arg	Gly
10		50					55					60				
	Ile	Ala	Glu	Ile	Thr	Glu	Met	Ile	His	Val	Ala	Ser	Leu	Leu	His	Asp
	65					70				75					80	
	Asp	Val	Leu	Asp	Asp	Ala	Asp	Thr	Arg	Arg	Gly	Val	Gly	Ser	Leu	Asn
					85					90					95	
15	Val	Val	Met	Gly	Asn	Lys	Val	Val	Ala	Leu	Leu	Ala	Thr	Ala	Val	Glu
				100					105					110		
	His	Leu	Val	Thr	Gly	Glu	Thr	Met	Glu	Ile	Thr	Ser	Ser	Thr	Glu	Gln
			115					120					125			
	Arg	Tyr	Ser	Met	Asp	Tyr	Tyr	Met	Gln	Lys	Thr	Tyr	Tyr	Lys	Thr	Ala
20		130					135					140				
	Ser	Leu	Ile	Ser	Asn	Ser	Cys	Lys	Ala	Val	Ala	Val	Leu	Thr	Gly	Gln
	145					150					155				160	
	Thr	Ala	Glu	Val	Ala	Val	Leu	Ala	Phe	Glu	Tyr	Gly	Arg	Asn	Leu	Gly
					165					170					175	
25	Leu	Ala	Phe	Gln	Leu	Ile	Asp	Asp	Ile	Leu	Asp	Phe	Thr	Gly	Thr	Ser
				180					185					190		
	Ala	Ser	Leu	Gly	Lys	Gly	Ser	Leu	Ser	Asp	Ile	Arg	His	Gly	Val	Ile
			195					200					205			
	Thr	Ala	Pro	Ile	Leu	Phe	Ala	Met	Glu	Glu	Phe	Pro	Gln	Leu	Arg	Glu
30		210					215					220				
	Val	Val	Asp	Gln	Val	Glu	Lys	Asp	Pro	Arg	Asn	Val	Asp	Ile	Ala	Leu
	225					230					235				240	
	Glu	Tyr	Leu	Gly	Lys	Ser	Lys	Gly	Ile	Gln	Arg	Ala	Arg	Glu	Leu	Ala
					245				250					255		
35	Met	Glu	His	Ala	Asn	Leu	Ala	Ala	Ala	Ala	Ile	Gly	Ser	Leu	Pro	Glu
				260				265						270		
	Thr	Asp	Asn	Glu	Asp	Val	Lys	Arg	Ser	Arg	Arg	Ala	Leu	Ile	Asp	Leu
			275					280					285			
	Thr	His	Arg	Val	Ile	Thr	Arg	Asn	Lys							
40		290					295									

&lt;210&gt; 10

&lt;211&gt; 561

&lt;212&gt; DNA

45 &lt;213&gt; Arabidopsis sp

&lt;400&gt; 10

	aagcgcatcc gtcctcttct acgattgccg ccagccgcat gtatggctgc ataaccgacc	60
	gcccctatcc gctcgcggcc gcggtcgaat tcattcacac cgcgacgctg ctgcatgacg	120
	acgtcgtcga tgaaagcgat ttgcgcgcgc gccgcgaaag cgcgcataag gttttcggca	180
5	atcaggcgag cgtgctcgtc ggcgatttcc ttttctcccg cgcttccag ctgatggtgg	240
	aagacggctc gctcgacgcg ctgcgcattc tctcgatgc ctccgccgtg atcgcgagg	300
	gcgaagtgat gcagctcggc accgcgcgca atcttgaaac caatatgagc cagtatctcg	360
	atgtgatcag cgcaagacc gccgcgctct ttgccgcgcg ctgcgaaatc ggcccgggtga	420
	tggcgaacgc gaaggcgga gatgctgccg cgatgtgcga atacggcatg aatctcggtg	480
10	tcgccttcca gatcatcgac gaccttctcg attacggcac cggcggccac gccgagcttg	540
	gcaagaacac gggcgacgat t	561

&lt;210&gt; 11

&lt;211&gt; 966

15 &lt;212&gt; DNA

&lt;213&gt; Arabidopsis sp

&lt;400&gt; 11

	atggtacttg ccgaggttcc aaagcttgcc tctgctgctg agtacttctt caaaaggggt	60
20	gtgcaaggaa aacagtttctg ttcaactatt ttgctgctga tggcgacagc tctgaatgta	120
	cgcgttccag aagcattgat tggggaatca acagatatag tcacatcaga attacgcgta	180
	aggcaacggg gtattgctga aatcactgaa atgatacacg tcgcaagtct actgcacgat	240
	gatgtcttgg atgatgccga tacaaggcgt ggtgttggtt ccttaaagtgt tgtaatgggt	300
	aacaagatgt cgggtattagc aggagacttc ttgctctccc gggcttggtg ggctctcgct	360
25	gcttttaaaga acacagaggt tgtagcatta cttgcaactg ctgtagaaca tcttgttacc	420
	ggtgaaacca tggaaataac tagttcaacc gagcagcgtt atagtatgga ctactacatg	480
	cagaagacat attataagac agcatcgcta atctctaaca gctgcaaagc tgttgccgtt	540
	ctcactggac aaacagcaga agttgccgtg ttagcttttg agtatgggag gaatctgggt	600
	ttagcattcc aattaataga cgacattctt gatttcacgg gcacatctgc ctctctcgga	660
30	aagggatcgt tgtcagatat tcgccatgga gtcataacag cccaatcct ctttgccatg	720
	gaagagtttc ctcaactacg cgaagttgtt gatcaagttg aaaaagatcc taggaatgtt	780
	gacattgctt tagagtatct tgggaagagc aagggaatac agaggggcaag agaattagcc	840
	atggaacatg cgaatctagc agcagctgca atcgggtctc tacctgaaac agacaatgaa	900
	gatgtcaaaa gatcgaggcg ggcacttatt gacttgaccc atagagtcac caccagaaac	960
35	aagtga	966

&lt;210&gt; 12

&lt;211&gt; 321

&lt;212&gt; PRT

40 &lt;213&gt; Arabidopsis sp

&lt;400&gt; 12

	Met Val Leu Ala Glu Val Pro Lys Leu Ala Ser Ala Ala Glu Tyr Phe	
	1                    5                    10                    15	
45	Phe Lys Arg Gly Val Gln Gly Lys Gln Phe Arg Ser Thr Ile Leu Leu	
	20                    25                    30	

	Leu	Met	Ala	Thr	Ala	Leu	Asn	Val	Arg	Val	Pro	Glu	Ala	Leu	Ile	Gly	
5	Ile	Ala	Glu	Ile	Thr	Glu	Met	Ile	His	Val	Ala	Ser	Leu	Leu	His	Asp	
10	Val	Val	Met	Gly	Asn	Lys	Met	Ser	Val	Leu	Ala	Gly	Asp	Phe	Leu	Leu	
15	Glu	Ile	Thr	Ser	Ser	Thr	Glu	Gln	Arg	Tyr	Ser	Met	Asp	Tyr	Tyr	Met	
20	Ala	Val	Ala	Val	Leu	Thr	Gly	Gln	Thr	Ala	Glu	Val	Ala	Val	Leu	Ala	
25	Ser	Asp	Ile	Arg	His	Gly	Val	Ile	Thr	Ala	Pro	Ile	Leu	Phe	Ala	Met	
30	Pro	Arg	Asn	Val	Asp	Ile	Ala	Leu	Glu	Tyr	Leu	Gly	Lys	Ser	Lys	Gly	
35	Ser	Arg	Arg	Ala	Leu	Ile	Asp	Leu	Thr	His	Arg	Val	Ile	Thr	Arg	Asn	
40	<210>	13															
	<211>	621															
	<212>	DNA															
	<213>	Arabidopsis sp															
45	<400>	13															
	gctttctctt	ttgctaattc	ttgagctttc	ttgatccac	cgcgatttct	aactatttca											60

	atcgcttctt	caagcgatcc	aggctcacia	aactcagact	caatgatctc	tcttagcctt	120
	ggctcattct	ctagecgcaa	gatcactggc	gccgttatgt	tacctttggc	taagtcatta	180
	gctgcaggct	tacctaactg	ctctgtggac	tgagtgaagt	ccagaatgtc	atcaactact	240
	tgaaaagata	aaccgagatt	cttcccgaac	tgatacattt	gctctgcgac	cttgctttcg	300
5	actttactga	aaattgctgc	tcctttgggtg	cttgcagcta	ctaataaagc	tgtctttag	360
	taactcttta	gcatgtagtc	atcaagcttg	acatcacaat	cgaataaact	cgatgcttgc	420
	tttatctcac	cgcttgcaaa	atctttgatc	acctgcaaaa	agataaatca	agattcagac	480
	caaatgttct	ttgtattgag	tagcttcac	taatctcaga	aaggaatatt	acctgactta	540
	tgagcttaat	gacttcaagg	ttttcgagat	ttgtaagtac	catgatgctt	gagcaacatg	600
10	aaatccccag	ctaatacagc	t				621

&lt;210&gt; 14

&lt;211&gt; 741

&lt;212&gt; DNA

15 &lt;213&gt; Arabidopsis sp

&lt;400&gt; 14

	ggtgagtttt	gttaatagtt	atgagattca	tctatTTTTg	tcataaaatt	gtttggtttg	60
	gtttaaactc	tgtgtataat	tgcaggaaag	gaaacagttc	atgagctttt	cggcacaaga	120
20	gtagcgggtg	tagctggaga	tttcatgttt	gctcaagcgt	catggtactt	agcaaactct	180
	gagaatcttg	aagttattaa	gctcatcagt	caggtactta	gttactctta	cattgttttt	240
	ctatgagggt	gagctatgaa	tctcattttc	ttgaataatg	ctgtgcctca	aacttttttt	300
	catgtttttc	ggtgatcaaa	gactttgcaa	gcggagagat	aaagcaggcg	tccagcttat	360
	ttgactgcga	caccaagctc	gacgagtact	tactcaaaag	tttctacaag	acagcctctt	420
25	tagtggctgc	gagcaccaaa	ggagctgcc	ttttcagcag	agttgagcct	gatgtgacag	480
	aacaaatgta	cgagtttggg	aagaatctcg	gtctctcttt	ccagatagtt	gatgatattt	540
	tggatttcac	tcagtcgaca	gagcagctcg	ggaagccagc	agggagtgat	ttggctaaag	600
	gtaacttaac	agcacctgtg	attttcgctc	tggagagggg	gccaaaggct	agagagatca	660
	ttgagtcaaa	gttctgtgag	gcgggttctc	tggagaagc	gattgaagcg	gtgacaaaag	720
30	gtggggggat	taagagagca	c				741

&lt;210&gt; 15

&lt;211&gt; 1087

&lt;212&gt; DNA

35 &lt;213&gt; Arabidopsis sp

&lt;400&gt; 15

	cctcttcagc	caatccagag	gaagaagaga	caacttttta	tctttcgtca	agagtctccg	60
	aaaacgcacg	gttttatgct	ctctcttctg	ccctcacctc	acaagacgca	gggcacatga	120
40	ttcaaccaga	gggaaaaagc	aacgataaca	actctgcttt	tgatttcaag	ctgtatatga	180
	tccgcaaagc	cgagtctgta	aatgcggctc	tcgacgtttc	cgtaccgctt	ctgaaacccc	240
	ttacgatcca	agaagcggtc	aggtactctt	tgctagccgg	cggaaaacgt	gtgaggcctc	300
	tgctctgcat	tgccgcttgt	gagcttgtgg	ggggcgacga	ggctactgcc	atgtcagccg	360
	cttgccggt	cgagatgatc	cacacaagct	ctctcattca	tgacgatctt	ccgtgcatgg	420
45	acaatgccga	cctccgtaga	ggcaagccca	ccaatcacia	ggtatgttgt	ttaattatat	480
	gaaggctcag	agataatgct	gaactagtgt	tgaaccaatt	tttgctcaaa	caaggtatat	540

ggagaagaca tggcgggtttt ggcaggtgat gcactccttg cattggcgtt tgagcacatg 600  
 acggttgtgt cgagtgggtt ggtcgctccc gagaagatga ttgcgcgcgt gggtgagctg 660  
 gccagggcca tagggactac agggctagtt gctggacaaa tgatagacct agccagcgaa 720  
 agactgaatc cagacaaggt tggattggag catctagagt tcatccatct ccacaaaacg 780  
 5 gcggcattgt tggaggcagc ggcagtttta ggggttataa tgggaggttg aacagaggaa 840  
 gaaatcgaaa agcttagaaa gtatgctagg tgtattggac tactgtttca gggtgttgat 900  
 gacattctcg acgtaacaaa atctactgag gaattgggta agacagccgg aaaagacgta 960  
 atggccggaa agctgacgta tccaaggctg ataggtttgg agggatccag ggaagttgca 1020  
 gagcacctga ggagagaagc agaggaaaag cttaaagggt ttgatccaag tcaggcggcg 1080  
 10 cctctgg 1087

<210> 16

<211> 1164

<212> DNA

15 <213> Arabidopsis sp

<400> 16

atgacttcga ttctcaacac tgtctccacc atccactctt ccagagttac ctccgtcgat 60  
 cgagtcggag tctctctctt tcggaattcg gattccgttg agttcactcg ccggcgttct 120  
 20 ggtttctcga cgttgatcta cgaatcaccg gggcggagat ttgttggtcg tgcggcggag 180  
 actgatactg ataaagttaa atctcagaca cctgacaagg caccagccgg tggttcaagc 240  
 attaaccagc ttctcggtat caaaggagca tctcaagaaa ctaataaatg gaagattcgt 300  
 cttcagctta caaaaccagt cacttggcct ccactgggtt ggggagtcgt ctgtggtgct 360  
 gctgcttcag ggaactttca ttggacccca gaggatgttg ctaagtcgat tctttgcatg 420  
 25 atgatgtctg gtcttgtct tactggctat acacagacaa tcaacgactg gtatgataga 480  
 gatatcgacg caattaatga gccatatcgt ccaattccat ctggagcaat atcagagcca 540  
 gaggttatta cacaagtctg ggtgctatta ttgggaggtc ttggtattgc tggaaatatta 600  
 gatgtgtggg cagggcatac cactcccact gtcttctatc ttgctttggg aggatcattg 660  
 ctatcttata tatactctgc tccacctctt aagctaaaac aaaatggatg gggtggaaat 720  
 30 tttgcacttg gagcaagcta tattagtttg ccatgggtgg ctggccaagc attgtttggc 780  
 actcttacgc cagatgttgt tgttctaaca ctcttgta gcatagctgg gttaggaata 840  
 gccattgta acgacttcaa aagtgttgaa ggagatagag cattaggact tcagtctctc 900  
 ccagtagctt ttggeaccga aactgcaaaa tggatatgcg ttggtgctat agacattact 960  
 cagctttctg ttgccggata tctattagca tctgggaaac cttattatgc gttggcgttg 1020  
 35 gttgctttga tcattcctca gattgtgttc cagtttaaact actttctcaa ggaccctgtc 1080  
 aaatacgacg tcaagtacca ggcaagcgcg cagccattct tgggtgctcg aatatttgta 1140  
 acggcattag catcgcaaca ctga 1164

<210> 17

40 <211> 387

<212> PRT

<213> Arabidopsis sp

<400> 17

45 Met Thr Ser Ile Leu Asn Thr Val Ser Thr Ile His Ser Ser Arg Val

1

5

10

15



	Thr	Ser	Val	Asp	Arg	Val	Gly	Val	Leu	Ser	Leu	Arg	Asn	Ser	Asp	Ser	
				20					25					30			
	Val	Glu	Phe	Thr	Arg	Arg	Arg	Ser	Gly	Phe	Ser	Thr	Leu	Ile	Tyr	Glu	
			35					40					45				
5	Ser	Pro	Gly	Arg	Arg	Phe	Val	Val	Arg	Ala	Ala	Glu	Thr	Asp	Thr	Asp	
		50					55					60					
	Lys	Val	Lys	Ser	Gln	Thr	Pro	Asp	Lys	Ala	Pro	Ala	Gly	Gly	Ser	Ser	
	65					70					75					80	
	Ile	Asn	Gln	Leu	Leu	Gly	Ile	Lys	Gly	Ala	Ser	Gln	Glu	Thr	Asn	Lys	
10					85					90					95		
	Trp	Lys	Ile	Arg	Leu	Gln	Leu	Thr	Lys	Pro	Val	Thr	Trp	Pro	Pro	Leu	
				100					105					110			
	Val	Trp	Gly	Val	Val	Cys	Gly	Ala	Ala	Ala	Ser	Gly	Asn	Phe	His	Trp	
			115					120					125				
15	Thr	Pro	Glu	Asp	Val	Ala	Lys	Ser	Ile	Leu	Cys	Met	Met	Met	Ser	Gly	
			130				135					140					
	Pro	Cys	Leu	Thr	Gly	Tyr	Thr	Gln	Thr	Ile	Asn	Asp	Trp	Tyr	Asp	Arg	
	145					150					155					160	
	Asp	Ile	Asp	Ala	Ile	Asn	Glu	Pro	Tyr	Arg	Pro	Ile	Pro	Ser	Gly	Ala	
20					165					170					175		
	Ile	Ser	Glu	Pro	Glu	Val	Ile	Thr	Gln	Val	Trp	Val	Leu	Leu	Leu	Gly	
				180					185				190				
	Gly	Leu	Gly	Ile	Ala	Gly	Ile	Leu	Asp	Val	Trp	Ala	Gly	His	Thr	Thr	
			195					200				205					
25	Pro	Thr	Val	Phe	Tyr	Leu	Ala	Leu	Gly	Gly	Ser	Leu	Leu	Ser	Tyr	Ile	
		210					215					220					
	Tyr	Ser	Ala	Pro	Pro	Leu	Lys	Leu	Lys	Gln	Asn	Gly	Trp	Val	Gly	Asn	
	225					230					235					240	
	Phe	Ala	Leu	Gly	Ala	Ser	Tyr	Ile	Ser	Leu	Pro	Trp	Trp	Ala	Gly	Gln	
30					245					250					255		
	Ala	Leu	Phe	Gly	Thr	Leu	Thr	Pro	Asp	Val	Val	Val	Leu	Thr	Leu	Leu	
			260					265					270				
	Tyr	Ser	Ile	Ala	Gly	Leu	Gly	Ile	Ala	Ile	Val	Asn	Asp	Phe	Lys	Ser	
		275						280				285					
35	Val	Glu	Gly	Asp	Arg	Ala	Leu	Gly	Leu	Gln	Ser	Leu	Pro	Val	Ala	Phe	
		290					295					300					
	Gly	Thr	Glu	Thr	Ala	Lys	Trp	Ile	Cys	Val	Gly	Ala	Ile	Asp	Ile	Thr	
	305					310					315					320	
	Gln	Leu	Ser	Val	Ala	Gly	Tyr	Leu	Leu	Ala	Ser	Gly	Lys	Pro	Tyr	Tyr	
40					325					330					335		
	Ala	Leu	Ala	Leu	Val	Ala	Leu	Ile	Ile	Pro	Gln	Ile	Val	Phe	Gln	Phe	
			340					345					350				
	Lys	Tyr	Phe	Leu	Lys	Asp	Pro	Val	Lys	Tyr	Asp	Val	Lys	Tyr	Gln	Ala	
		355						360				365					
45	Ser	Ala	Gln	Pro	Phe	Leu	Val	Leu	Gly	Ile	Phe	Val	Thr	Ala	Leu	Ala	
		370					375					380					

Ser Gln His

385

&lt;210&gt; 18

5 &lt;211&gt; 981

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis sp

&lt;400&gt; 18

10	atgttggttta	gtgggttcagc	gatcccatta	agcagcttct	gctctcttcc	ggagaaaccc	60
	cacactcttc	ctatgaaact	ctctcccgt	gcaatccgat	cttcatectc	atctgccccg	120
	gggtcggtga	acttcgatct	gaggacgtat	tggacgactc	tgatcaccga	gatcaaccag	180
	aagctggatg	aggccatacc	ggtcaagcac	cctgcgggga	tctacgaggc	tatgagatac	240
	tctgtactcg	cacaaggcgc	caagcgtgcc	cctcctgtga	tgtgtgtggc	ggcctgcgag	300
15	ctcttcgggtg	gcgategcct	cgccgcttcc	cccaccgcct	gtgccctaga	aatgggtgcac	360
	gcggcttcgt	tgatacacga	cgacctcccc	tgtatggacg	acgatcctgt	gcgcagagga	420
	aagccatcta	accacactgt	ctacggctct	ggcatggcca	ttctcgccgg	tgacgccctc	480
	ttcccactcg	ccttccagca	cattgtctcc	cacacgcctc	ctgaccttgt	tccccgagcc	540
	accatectca	gactcatcac	tgagattgcc	cgcactgtcg	gctccactgg	tatggctgca	600
20	ggccagtacg	tcgaccttga	aggagggtccc	tttcctcttt	cctttgttca	ggagaagaaa	660
	ttcggagcca	tgggtgaatg	ctctgccgtg	tgcgggtggc	tattgggcgg	tgccactgag	720
	gatgagctcc	agagtctccg	aagggtacggg	agagccgtcg	ggatgctgta	tcagggtggc	780
	gatgacatca	ccgaggacaa	gaagaagagc	tatgatgggtg	gagcagagaa	gggaatgatg	840
	gaaatggcgg	aagagctcaa	ggagaaggcg	aagaaggagc	ttcaagtgtt	tgacaacaag	900
25	tatggaggag	gagacacact	tgttcctctc	tacaccttcg	ttgactacgc	tgctcatcga	960
	cattttcttc	ttcccctctg	a				981

&lt;210&gt; 19

&lt;211&gt; 245

30 &lt;212&gt; DNA

&lt;213&gt; Glycine sp

&lt;400&gt; 19

	gcaacatctg	ggactggggtt	tgtcttggggg	agtggtagtg	ctgttgatct	ttcggcactt	60
35	tcttgcaactt	gcttgggtac	catgatgggtt	gctgcatctg	ctaactcttt	gaatcaggtg	120
	tttgagatca	ataatgatgc	taaaatgaag	agaacaagtc	gcaggccact	accctcagga	180
	cgcatacaca	tacctcatgc	agttggctgg	gcatacctctg	ttggattagc	tggtacggct	240
	ctact						245

40 &lt;210&gt; 20

&lt;211&gt; 253

&lt;212&gt; DNA

&lt;213&gt; Glycine sp

45 &lt;400&gt; 20

	attggcttcc	caagatcatt	gggttttctt	gttgcatcca	tgaccttcta	ctccttgggt	60
--	------------	------------	------------	------------	------------	------------	----

ttggcattgt ccaaggatat acctgacgtt gaaggagata aagagcacgg cattgattct 120  
 tttgcagtac gtctaggtca gaaacgggca ttttggattt gcgtttcctt ttttgaaatg 180  
 gctttcggag ttggtatcct ggccggagca tcatgctcac acttttggac taaaattttc 240  
 acgggtatgg gaa 253

5

<210> 21  
 <211> 275  
 <212> DNA  
 <213> Glycine sp

10

<400> 21  
 tgatcttcta ctctctgggt atggcattgt ccaaggatat atctgacgtt aaaggagata 60  
 aagcatacgg catcgatact ttagcgatac gtttgggtca aaaatgggta ttttggattt 120  
 gcattatcct ttttgaaatg gcttttggag ttgccctctt ggcaggagca acatcttctt 180  
 15 acctttggat taaaattgtc acgggtctgg gacatgctat tcttgcttca attctcttgt 240  
 accaagccaa atctatatac ttgagcaaca aagtt 275

<210> 22  
 <211> 299  
 20 <212> DNA  
 <213> Glycine sp

<220>  
 <221> misc\_feature  
 25 <222> (1) ... (299)  
 <223> n = A,T,C or G

<400> 22  
 ccanaatang tncatcttng aaagacaatt ggccctcttca acacacaagt ctgcatgtga 60  
 30 agaagaggcc aattgtcttt ccaagatcac ttatngtggc tattgtaatc atgaacttct 120  
 tctttgtggg tatggcattg gcaaaggata tacctanctg ttgaaggaga taaaatatat 180  
 ggcattgata cttttgcaat acgtataggt caaaaacaag tatttttggat ttgtattttc 240  
 ctttttgaaa ggctttcgga gtttccctag tggcaggagc aacatcttct agccttggg 299

35 <210> 23  
 <211> 767  
 <212> DNA  
 <213> Glycine sp

40 <400> 23  
 gtggaggctg tgggtgctgc cctgtttatg aatattttata ttgttggttt gaatcaattg 60  
 tctgatgttg aaatagacaa gataaacaag ccgtatcttc cattagcatc tggggaatat 120  
 tcctttgaaa ctgggtgtcac tattgttgca tctttttcaa ttctgagttt ttggcttggc 180  
 tgggttgtag gttcatggcc attattttgg gccctttttg taagctttgt gctaggaact 240  
 45 gcttattcaa tcaatgtgcc tctgttgaga tggaagaggt ttgcagtgtc tgcagcgatg 300  
 tgcattctag ctgttcgggc agtaatagtt caacttgcac ttttccctca catgcagact 360

catgtgtaca agaggccacc tgtctttttca agaccattga tttttgctac tgcattcatg 420  
 agcttcttct ctgtagttat agcactgttt aaggatatac ctgacattga aggagataaa 480  
 gtatttggca tccaatcttt ttcagtgtgt ttaggtcaga agccgggtgt ctggacttgt 540  
 gttacccttc ttgaaatagc ttatggagtc gccctcctgg tgggagctgc atctccttgt 600  
 5 ctttggagca aaattttcac gggctctggga cacgctgtgc tggcttcaat tctctgggtt 660  
 catgccaaat ctgtagattt gaaaagcaaa gcttcgataa catccttcta tatgtttatt 720  
 tggaagctat tttatgcaga atacttactc attccttttg ttagatg 767

<210> 24

10 <211> 255

<212> PRT

<213> Glycine sp

<400> 24

15 Val Glu Ala Val Val Ala Ala Leu Phe Met Asn Ile Tyr Ile Val Gly  
 1 5 10 15  
 Leu Asn Gln Leu Ser Asp Val Glu Ile Asp Lys Ile Asn Lys Pro Tyr  
 20 25 30  
 Leu Pro Leu Ala Ser Gly Glu Tyr Ser Phe Glu Thr Gly Val Thr Ile  
 35 40 45  
 20 Val Ala Ser Phe Ser Ile Leu Ser Phe Trp Leu Gly Trp Val Val Gly  
 50 55 60  
 Ser Trp Pro Leu Phe Trp Ala Leu Phe Val Ser Phe Val Leu Gly Thr  
 65 70 75 80  
 25 Ala Tyr Ser Ile Asn Val Pro Leu Leu Arg Trp Lys Arg Phe Ala Val  
 85 90 95  
 Leu Ala Ala Met Cys Ile Leu Ala Val Arg Ala Val Ile Val Gln Leu  
 100 105 110  
 Ala Phe Phe Leu His Met Gln Thr His Val Tyr Lys Arg Pro Pro Val  
 115 120 125  
 30 Phe Ser Arg Pro Leu Ile Phe Ala Thr Ala Phe Met Ser Phe Phe Ser  
 130 135 140  
 Val Val Ile Ala Leu Phe Lys Asp Ile Pro Asp Ile Glu Gly Asp Lys  
 145 150 155 160  
 35 Val Phe Gly Ile Gln Ser Phe Ser Val Cys Leu Gly Gln Lys Pro Val  
 165 170 175  
 Phe Trp Thr Cys Val Thr Leu Leu Glu Ile Ala Tyr Gly Val Ala Leu  
 180 185 190  
 Leu Val Gly Ala Ala Ser Pro Cys Leu Trp Ser Lys Ile Phe Thr Gly  
 195 200 205  
 40 Leu Gly His Ala Val Leu Ala Ser Ile Leu Trp Phe His Ala Lys Ser  
 210 215 220  
 Val Asp Leu Lys Ser Lys Ala Ser Ile Thr Ser Phe Tyr Met Phe Ile  
 225 230 235 240  
 45 Trp Lys Leu Phe Tyr Ala Glu Tyr Leu Leu Ile Pro Phe Val Arg  
 245 250 255

<210> 25  
 <211> 360  
 <212> DNA  
 5 <213> Zea sp  
  
 <220>  
 <221> misc\_feature  
 <222> (1)...(360)  
 10 <223> n = A,T,C or G  
  
 <400> 25  
 ggcgtcttca cttgtttctgg tcttctcgta tcccctgatg aagaggttca cattttggcc 60  
 tcaggccttat cttggcctga cattcaactg gggagcttta ctagggtggg ctgctattaa 120  
 15 ggaaagcata gaccctgcaa atcatccttc cattgtatac agctgggtatt tggtggacgc 180  
 tgggtgtatga tactatatat gcgcacacagg tgtttcgcta tccctacttt catattaatc 240  
 cttgatgaag tggccatttc atgttgctgc ggtgggtctta tacttgcata tctccatgca 300  
 tctcaggaca aagangatga cctgaaagta ggagtccaag tccacagctt aagatttggg 360  
  
 20 <210> 26  
 <211> 299  
 <212> DNA  
 <213> Zea sp  
  
 25 <220>  
 <221> misc\_feature  
 <222> (1)...(299)  
 <223> n = A,T,C or G  
  
 30 <400> 26  
 gatggttgca gcatctgcaa ataccctcaa ccagggtgttt gngataaaaa atgatgctaa 60  
 aatgaaaagg acaatgcgtg cccctgcca tctggctgca ttagtcctgc acatgctgcg 120  
 atgtgggcta caagtgttgg agttgcagga acagctttgt tggcctggaa ggctaattggc 180  
 ttggcagctg ggcttgacgc ttctaattctt gttctgtatg catttgtgta tacgccgttg 240  
 35 aagcaaatac accctgttaa tacatgggtt ggggcagtcg ttggtgccat cccaccact 299  
  
 <210> 27  
 <211> 255  
 <212> DNA  
 40 <213> Zea sp  
  
 <220>  
 <221> misc\_feature  
 <222> (1)...(255)  
 45 <223> n = A,T,C or G

&lt;400&gt; 27

anacttgc atctccatgc ntctcaggac aaagangatg acctgaaagt aggtgtcaag 60  
 tccacagcat taagatttgg agatttgacc nnatactgna tcagtggctt tggcgcgga 120  
 tgcttcggca gcttagcact cagtgggttac aatgctgacc ttggttggtg tttagtgtga 180  
 5 tgcttgagcg aagaatggta tngtttttac ttgatattga ctccagacct gaaatcatgt 240  
 tggacagggt ggccc 255

&lt;210&gt; 28

&lt;211&gt; 257

10 &lt;212&gt; DNA

&lt;213&gt; Zea sp

&lt;400&gt; 28

attgaagggg ataggactct ggggcttcag tcacttcctg ttgcttttgg gatggaaact 60  
 15 gcaaaatgga tttgtgttgg agcaattgat atcactcaat tatctgttgc aggttaccta 120  
 ttgagcaccg gtaagctgta ttatgccctg gtgttgcttg ggctaacaat tcctcagggtg 180  
 ttctttcagt tccagtactt cctgaaggac cctgtgaagt atgatgtcaa atatcaggca 240  
 agcgcacaa cattctt 257

20 &lt;210&gt; 29

&lt;211&gt; 368

&lt;212&gt; DNA

&lt;213&gt; Zea sp

25 &lt;400&gt; 29

atccagttgc aaataataat ggcggttcttc tctgttgtaa tagcactatt caaggatata 60  
 cctgacatcg aaggggaccg catattcggg atccgatact tcagcgctcg gttagggcaa 120  
 aagaaggtct tttggatctg cgttggcttg cttgagatgg cctacagcgt tgcgatactg 180  
 atgggagcta cctcttcctg tttgtggagc aaaacagcaa ccatcgctgg ccattccata 240  
 30 cttgccgcga tcctatggag ctgcgcgcga tcggtggact tgacgagcaa agccgcaata 300  
 acgtccttct acatgttcat ctggaagctg ttctacgcgg agtacctgct catccctctg 360  
 gtgcggtg 368

&lt;210&gt; 30

35 &lt;211&gt; 122

&lt;212&gt; PRT

&lt;213&gt; Zea sp

&lt;400&gt; 30

40 Ile Gln Leu Gln Ile Ile Met Ala Phe Phe Ser Val Val Ile Ala Leu  
 1 5 10 15  
 Phe Lys Asp Ile Pro Asp Ile Glu Gly Asp Arg Ile Phe Gly Ile Arg  
 20 25 30  
 Ser Phe Ser Val Arg Leu Gly Gln Lys Lys Val Phe Trp Ile Cys Val  
 45 35 40 45  
 Gly Leu Leu Glu Met Ala Tyr Ser Val Ala Ile Leu Met Gly Ala Thr

50 55 60  
 Ser Ser Cys Leu Trp Ser Lys Thr Ala Thr Ile Ala Gly His Ser Ile  
 65 70 75 80  
 Leu Ala Ala Ile Leu Trp Ser Cys Ala Arg Ser Val Asp Leu Thr Ser  
 5 85 90 95  
 Lys Ala Ala Ile Thr Ser Phe Tyr Met Phe Ile Trp Lys Leu Phe Tyr  
 100 105 110  
 Ala Glu Tyr Leu Leu Ile Pro Leu Val Arg  
 115 120  
 10  
 <210> 31  
 <211> 278  
 <212> DNA  
 <213> Zea sp  
 15  
 <400> 31  
 tattcagcac cacctctcaa gctcaagcag aatggatgga ttgggaactt cgctctgggt 60  
 gcgagttaca tcagcttgcc ctggtgggct ggccaggcgt tatttggaac tcttacacca 120  
 gatattcattg tcttgactac tttgtacagc atagctgggc tagggattgc tattgtaaat 180  
 20 gatttcaaga gtattgaagg ggataggact ctggggcttc agtcacttcc tgttgctttt 240  
 gggatggaaa ctgcaaaatg gatttgtgtt ggagcaat 278  
  
 <210> 32  
 <211> 292  
 25 <212> PRT  
 <213> Synechocystis sp  
  
 <400> 32  
 Met Val Ala Gln Thr Pro Ser Ser Pro Pro Leu Trp Leu Thr Ile Ile  
 30 1 5 10 15  
 Tyr Leu Leu Arg Trp His Lys Pro Ala Gly Arg Leu Ile Leu Met Ile  
 20 25 30  
 Pro Ala Leu Trp Ala Val Cys Leu Ala Ala Gln Gly Leu Pro Pro Leu  
 35 35 40 45  
 Pro Leu Leu Gly Thr Ile Ala Leu Gly Thr Leu Ala Thr Ser Gly Leu  
 50 55 60  
 Gly Cys Val Val Asn Asp Leu Trp Asp Arg Asp Ile Asp Pro Gln Val  
 65 70 75 80  
 Glu Arg Thr Lys Gln Arg Pro Leu Ala Ala Arg Ala Leu Ser Val Gln  
 40 85 90 95  
 Val Gly Ile Gly Val Ala Leu Val Ala Leu Leu Cys Ala Ala Gly Leu  
 100 105 110  
 Ala Phe Tyr Leu Thr Pro Leu Ser Phe Trp Leu Cys Val Ala Ala Val  
 115 120 125  
 45 Pro Val Ile Val Ala Tyr Pro Gly Ala Lys Arg Val Phe Pro Val Pro  
 130 135 140



Gln Leu Val Leu Ser Ile Ala Trp Gly Phe Ala Val Leu Ile Ser Trp  
 145 150 155 160  
 Ser Ala Val Thr Gly Asp Leu Thr Asp Ala Thr Trp Val Leu Trp Gly  
 165 170 175  
 5 Ala Thr Val Phe Trp Thr Leu Gly Phe Asp Thr Val Tyr Ala Met Ala  
 180 185 190  
 Asp Arg Glu Asp Asp Arg Arg Ile Gly Val Asn Ser Ser Ala Leu Phe  
 195 200 205  
 Phe Gly Gln Tyr Val Gly Glu Ala Val Gly Ile Phe Phe Ala Leu Thr  
 10 210 215 220  
 Ile Gly Cys Leu Phe Tyr Leu Gly Met Ile Leu Met Leu Asn Pro Leu  
 225 230 235 240  
 Tyr Trp Leu Ser Leu Ala Ile Ala Ile Val Gly Trp Val Ile Gln Tyr  
 245 250 255  
 15 Ile Gln Leu Ser Ala Pro Thr Pro Glu Pro Lys Leu Tyr Gly Gln Ile  
 260 265 270  
 Phe Gly Gln Asn Val Ile Ile Gly Phe Val Leu Leu Ala Gly Met Leu  
 275 280 285  
 Leu Gly Trp Leu  
 20 290  
  
 <210> 33  
 <211> 316  
 <212> PRT  
 25 <213> Synechocystis sp  
  
 <400> 33  
 Met Val Thr Ser Thr Lys Ile His Arg Gln His Asp Ser Met Gly Ala  
 1 5 10 15  
 30 Val Cys Lys Ser Tyr Tyr Gln Leu Thr Lys Pro Arg Ile Ile Pro Leu  
 20 25 30  
 Leu Leu Ile Thr Thr Ala Ala Ser Met Trp Ile Ala Ser Glu Gly Arg  
 35 40 45  
 Val Asp Leu Pro Lys Leu Leu Ile Thr Leu Leu Gly Gly Thr Leu Ala  
 50 55 60  
 35 Ala Ala Ser Ala Gln Thr Leu Asn Cys Ile Tyr Asp Gln Asp Ile Asp  
 65 70 75 80  
 Tyr Glu Met Leu Arg Thr Arg Ala Arg Pro Ile Pro Ala Gly Lys Val  
 85 90 95  
 40 Gln Pro Arg His Ala Leu Ile Phe Ala Leu Ala Leu Gly Val Leu Ser  
 100 105 110  
 Phe Ala Leu Leu Ala Thr Phe Val Asn Val Leu Ser Gly Cys Leu Ala  
 115 120 125  
 Leu Ser Gly Ile Val Phe Tyr Met Leu Val Tyr Thr His Trp Leu Lys  
 45 130 135 140  
 Arg His Thr Ala Gln Asn Ile Val Ile Gly Gly Ala Ala Gly Ser Ile

	145		150		155		160									
	Pro	Pro	Leu	Val	Gly	Trp	Ala	Ala	Val	Thr	Gly	Asp	Leu	Ser	Trp	Thr
					165					170					175	
5	Pro	Trp	Val	Leu	Phe	Ala	Leu	Ile	Phe	Leu	Trp	Thr	Pro	Pro	His	Phe
				180					185					190		
	Trp	Ala	Leu	Ala	Leu	Met	Ile	Lys	Asp	Asp	Tyr	Ala	Gln	Val	Asn	Val
			195					200					205			
	Pro	Met	Leu	Pro	Val	Ile	Ala	Gly	Glu	Glu	Lys	Thr	Val	Ser	Gln	Ile
		210					215					220				
10	Trp	Tyr	Tyr	Ser	Leu	Leu	Val	Val	Pro	Phe	Ser	Leu	Leu	Leu	Val	Tyr
		225				230					235				240	
	Pro	Leu	His	Gln	Leu	Gly	Ile	Leu	Tyr	Leu	Ala	Ile	Ala	Ile	Ile	Leu
				245						250					255	
	Gly	Gly	Gln	Phe	Leu	Val	Lys	Ala	Trp	Gln	Leu	Lys	Gln	Ala	Pro	Gly
15				260					265					270		
	Asp	Arg	Asp	Leu	Ala	Arg	Gly	Leu	Phe	Lys	Phe	Ser	Ile	Phe	Tyr	Leu
			275					280					285			
	Met	Leu	Leu	Cys	Leu	Ala	Met	Val	Ile	Asp	Ser	Leu	Pro	Val	Thr	His
		290					295					300				
20	Gln	Leu	Val	Ala	Gln	Met	Gly	Thr	Leu	Leu	Leu	Gly				
	305					310					315					

&lt;210&gt; 34

&lt;211&gt; 324

25 &lt;212&gt; PRT

&lt;213&gt; Synechocystis sp

&lt;400&gt; 34

	Met	Ser	Asp	Thr	Gln	Asn	Thr	Gly	Gln	Asn	Gln	Ala	Lys	Ala	Arg	Gln
30	1				5					10					15	
	Leu	Leu	Gly	Met	Lys	Gly	Ala	Ala	Pro	Gly	Glu	Ser	Ser	Ile	Trp	Lys
			20					25						30		
	Ile	Arg	Leu	Gln	Leu	Met	Lys	Pro	Ile	Thr	Trp	Ile	Pro	Leu	Ile	Trp
		35					40					45				
35	Gly	Val	Val	Cys	Gly	Ala	Ala	Ser	Ser	Gly	Gly	Tyr	Ile	Trp	Ser	Val
		50				55					60					
	Glu	Asp	Phe	Leu	Lys	Ala	Leu	Thr	Cys	Met	Leu	Leu	Ser	Gly	Pro	Leu
	65				70				75				80			
	Met	Thr	Gly	Tyr	Thr	Gln	Thr	Leu	Asn	Asp	Phe	Tyr	Asp	Arg	Asp	Ile
40				85				90					95			
	Asp	Ala	Ile	Asn	Glu	Pro	Tyr	Arg	Pro	Ile	Pro	Ser	Gly	Ala	Ile	Ser
			100					105					110			
	Val	Pro	Gln	Val	Val	Thr	Gln	Ile	Leu	Ile	Leu	Leu	Val	Ala	Gly	Ile
		115					120					125				
45	Gly	Val	Ala	Tyr	Gly	Leu	Asp	Val	Trp	Ala	Gln	His	Asp	Phe	Pro	Ile
	130					135					140					

Met Met Val Leu Thr Leu Gly Gly Ala Phe Val Ala Tyr Ile Tyr Ser  
 145 150 155 160  
 Ala Pro Pro Leu Lys Leu Lys Gln Asn Gly Trp Leu Gly Asn Tyr Ala  
 165 170 175  
 5 Leu Gly Ala Ser Tyr Ile Ala Leu Pro Trp Trp Ala Gly His Ala Leu  
 180 185 190  
 Phe Gly Thr Leu Asn Pro Thr Ile Met Val Leu Thr Leu Ile Tyr Ser  
 195 200 205  
 Leu Ala Gly Leu Gly Ile Ala Val Val Asn Asp Phe Lys Ser Val Glu  
 10 210 215 220  
 Gly Asp Arg Gln Leu Gly Leu Lys Ser Leu Pro Val Met Phe Gly Ile  
 225 230 235 240  
 Gly Thr Ala Ala Trp Ile Cys Val Ile Met Ile Asp Val Phe Gln Ala  
 245 250 255  
 15 Gly Ile Ala Gly Tyr Leu Ile Tyr Val His Gln Gln Leu Tyr Ala Thr  
 260 265 270  
 Ile Val Leu Leu Leu Leu Ile Pro Gln Ile Thr Phe Gln Asp Met Tyr  
 275 280 285  
 Phe Leu Arg Asn Pro Leu Glu Asn Asp Val Lys Tyr Gln Ala Ser Ala  
 20 290 295 300  
 Gln Pro Phe Leu Val Phe Gly Met Leu Ala Thr Gly Leu Ala Leu Gly  
 305 310 315 320  
 His Ala Gly Ile  
 25  
 <210> 35  
 <211> 307  
 <212> PRT  
 <213> Synechocystis sp  
 30  
 <400> 35  
 Met Thr Glu Ser Ser Pro Leu Ala Pro Ser Thr Ala Pro Ala Thr Arg  
 1 5 10 15  
 Lys Leu Trp Leu Ala Ala Ile Lys Pro Pro Met Tyr Thr Val Ala Val  
 35 20 25 30  
 Val Pro Ile Thr Val Gly Ser Ala Val Ala Tyr Gly Leu Thr Gly Gln  
 35 40 45  
 Trp His Gly Asp Val Phe Thr Ile Phe Leu Leu Ser Ala Ile Ala Ile  
 50 55 60  
 40 Ile Ala Trp Ile Asn Leu Ser Asn Asp Val Phe Asp Ser Asp Thr Gly  
 65 70 75 80  
 Ile Asp Val Arg Lys Ala His Ser Val Val Asn Leu Thr Gly Asn Arg  
 85 90 95  
 Asn Leu Val Phe Leu Ile Ser Asn Phe Phe Leu Leu Ala Gly Val Leu  
 45 100 105 110  
 Gly Leu Met Ser Met Ser Trp Arg Ala Gln Asp Trp Thr Val Leu Glu

	115		120		125
	Leu Ile Gly Val Ala Ile Phe Leu Gly Tyr Thr Tyr Gln Gly Pro Pro				
	130		135		140
5	Phe Arg Leu Gly Tyr Leu Gly Leu Gly Glu Leu Ile Cys Leu Ile Thr				
	145		150		155
	Phe Gly Pro Leu Ala Ile Ala Ala Tyr Tyr Ser Gln Ser Gln Ser				
		165		170	175
	Phe Ser Trp Asn Leu Leu Thr Pro Ser Val Phe Val Gly Ile Ser Thr				
		180		185	190
10	Ala Ile Ile Leu Phe Cys Ser His Phe His Gln Val Glu Asp Asp Leu				
		195		200	205
	Ala Ala Gly Lys Lys Ser Pro Ile Val Arg Leu Gly Thr Lys Leu Gly				
		210		215	220
	Ser Gln Val Leu Thr Leu Ser Val Val Ser Leu Tyr Leu Ile Thr Ala				
15	225		230		235
	Ile Gly Val Leu Cys His Gln Ala Pro Trp Gln Thr Leu Leu Ile Ile				
		245		250	255
	Ala Ser Leu Pro Trp Ala Val Gln Leu Ile Arg His Val Gly Gln Tyr				
		260		265	270
20	His Asp Gln Pro Glu Gln Val Ser Asn Cys Lys Phe Ile Ala Val Asn				
		275		280	285
	Leu His Phe Phe Ser Gly Met Leu Met Ala Ala Gly Tyr Gly Trp Ala				
		290		295	300
	Gly Leu Gly				

25 305

<210> 36

<211> 927

<212> DNA

30 <213> Synechocystis sp

<400> 36

	atggcaacta tccaagcttt ttggcgcttc tcccgccccc ataccatcat tgggtacaact	60
	ctgagcgtct gggctgtgta tctgttaact attctcgggg atggaaactc agttaactcc	120
35	cctgcttccc tggatttagt gttcggcgct tggctggcct gcctgttggg taatgtgtac	180
	attgtcggcc tcaaccaatt gtgggatgtg gacattgacc gcatcaataa gccgaatttg	240
	cccctagcta acggagattt ttctatcgcc cagggccggt ggattgtggg actttgtggc	300
	gttgcttccct tggcgatcgc ctggggatta gggctatggc tggggctaac ggtgggcatt	360
	agtttgatta ttggcacggc ctattcgggt cgcgcagtga ggtaaagcg cttttccctg	420
40	ctggcgggccc tgtgtattct gacgggtgcg ggaattgtgg ttaacttggg cttattttta	480
	tttttttagaa ttggtttagg ttatcccccc actttaataa ccccatctg ggttttgact	540
	ttatttatct tagttttcac cgtggcgatc gccattttta aagatgtgcc agatatggaa	600
	ggcgatcggc aatttaagat tcaaacttta actttgcaaa tcggcaaaca aaacgttttt	660
	cggggaacct taattttact cactggttgt tatttagcca tggcaatctg gggcttatgg	720
45	gcggctatgc ctttaaatac tgctttcttg attgtttccc atttgtgctt attagcetta	780
	ctctggtggc ggagtcgaga tgtacactta gaaagcaaaa ccgaaattgc tagtttttat	840

cagtttattt ggaagctatt tttcttagag tacttgctgt atcccttggc tctgtgggta 900  
 cctaattttt ctaatactat ttttttag 927

<210> 37

5 <211> 308

<212> PRT

<213> Synechocystis sp

<400> 37

10	Met	Ala	Thr	Ile	Gln	Ala	Phe	Trp	Arg	Phe	Ser	Arg	Pro	His	Thr	Ile
	1				5					10					15	
	Ile	Gly	Thr	Thr	Leu	Ser	Val	Trp	Ala	Val	Tyr	Leu	Leu	Thr	Ile	Leu
				20					25					30		
	Gly	Asp	Gly	Asn	Ser	Val	Asn	Ser	Pro	Ala	Ser	Leu	Asp	Leu	Val	Phe
15			35					40					45			
	Gly	Ala	Trp	Leu	Ala	Cys	Leu	Leu	Gly	Asn	Val	Tyr	Ile	Val	Gly	Leu
		50					55					60				
	Asn	Gln	Leu	Trp	Asp	Val	Asp	Ile	Asp	Arg	Ile	Asn	Lys	Pro	Asn	Leu
	65				70					75					80	
20	Pro	Leu	Ala	Asn	Gly	Asp	Phe	Ser	Ile	Ala	Gln	Gly	Arg	Trp	Ile	Val
				85						90					95	
	Gly	Leu	Cys	Gly	Val	Ala	Ser	Leu	Ala	Ile	Ala	Trp	Gly	Leu	Gly	Leu
				100					105					110		
	Trp	Leu	Gly	Leu	Thr	Val	Gly	Ile	Ser	Leu	Ile	Ile	Gly	Thr	Ala	Tyr
25			115					120					125			
	Ser	Val	Pro	Pro	Val	Arg	Leu	Lys	Arg	Phe	Ser	Leu	Leu	Ala	Ala	Leu
		130					135					140				
	Cys	Ile	Leu	Thr	Val	Arg	Gly	Ile	Val	Val	Asn	Leu	Gly	Leu	Phe	Leu
	145				150					155					160	
30	Phe	Phe	Arg	Ile	Gly	Leu	Gly	Tyr	Pro	Pro	Thr	Leu	Ile	Thr	Pro	Ile
				165					170					175		
	Trp	Val	Leu	Thr	Leu	Phe	Ile	Leu	Val	Phe	Thr	Val	Ala	Ile	Ala	Ile
				180					185					190		
	Phe	Lys	Asp	Val	Pro	Asp	Met	Glu	Gly	Asp	Arg	Gln	Phe	Lys	Ile	Gln
35			195					200					205			
	Thr	Leu	Thr	Leu	Gln	Ile	Gly	Lys	Gln	Asn	Val	Phe	Arg	Gly	Thr	Leu
		210					215					220				
	Ile	Leu	Leu	Thr	Gly	Cys	Tyr	Leu	Ala	Met	Ala	Ile	Trp	Gly	Leu	Trp
	225				230					235					240	
40	Ala	Ala	Met	Pro	Leu	Asn	Thr	Ala	Phe	Leu	Ile	Val	Ser	His	Leu	Cys
				245					250					255		
	Leu	Leu	Ala	Leu	Leu	Trp	Trp	Arg	Ser	Arg	Asp	Val	His	Leu	Glu	Ser
				260				265					270			
	Lys	Thr	Glu	Ile	Ala	Ser	Phe	Tyr	Gln	Phe	Ile	Trp	Lys	Leu	Phe	Phe
45			275					280					285			
	Leu	Glu	Tyr	Leu	Leu	Tyr	Pro	Leu	Ala	Leu	Trp	Leu	Pro	Asn	Phe	Ser

290 295 300

Asn Thr Ile Phe

305

5 <210> 38  
<211> 1092  
<212> DNA  
<213> Synechocystis sp

10 <400> 38  
atgaaatttc cgccccacag tggttaccat tggcaaggtc aatcaccttt ctttgaaggt 60  
tggtagctgc gcctgctttt gccccaatcc ggggaaagtt ttgcttttat gtactccatc 120  
gaaaatcctg ctagcgatca tcattacggc ggcggtgctg tgcaaatttt agggccggct 180  
acgaaaaaac aagaaaatca ggaagaccaa cttgtttggc ggacatttcc ctcggtaaaa 240  
15 aaattttggg ccagtcctcg ccagtttgcc ctagggcatt ggggaaaatg tagggataac 300  
aggcaggcga aaccctact ctccgaagaa ttttttgcca cgggtcaagga aggttatcaa 360  
atccatcaaa atcagcacca aggacaaatc attcatggcg atcgccattg tcgttggcag 420  
ttcacgtag aaccggaagt aacttggggg agtcctaacc gatttcctcg ggctacagcg 480  
ggttggcttt cttttttacc cttgtttgat ccggttggc aaattctttt agcccaaggt 540  
20 agagcgacg gctggctgaa atggcagagg gaacagtatg aatttgacca cgccctagtt 600  
tatgccgaaa aaaattgggg tctactcctt cctcccgtt ggttttggct ccaagcaaat 660  
tattttcctg accatccagg actgagcgtc actgccgctg gcggggaacg gattgttctt 720  
ggtcgccccg aagaggtagc tttaattggc ttacatcacc aaggtaattt ttacgaattt 780  
ggcccggggc atggcacagt cacttggcaa gtagctccct ggggcccgtt gcaattaaaa 840  
25 gccagcaatg ataggtattg ggtcaagttg tccggaaaaa cagataaaaa aggcagttta 900  
gtccacactc ccaccgcca gggcttacia ctcaactgcc gagataccac taggggctat 960  
ttgtatttgc aattgggac tgtgggtcac ggcctgatag tgcaagggga aacggacacc 1020  
gcggggctag aagttggagg tgattggggg ttaacagagg aaaatttgag caaaaaaaca 1080  
gtgccattct ga 1092

30 <210> 39  
<211> 363  
<212> PRT  
<213> Synechocystis sp

35 <400> 39  
Met Lys Phe Pro Pro His Ser Gly Tyr His Trp Gln Gly Gln Ser Pro  
1 5 10 15  
Phe Phe Glu Gly Trp Tyr Val Arg Leu Leu Leu Pro Gln Ser Gly Glu  
40 20 25 30  
Ser Phe Ala Phe Met Tyr Ser Ile Glu Asn Pro Ala Ser Asp His His  
35 40 45  
Tyr Gly Gly Gly Ala Val Gln Ile Leu Gly Pro Ala Thr Lys Lys Gln  
50 55 60  
45 Glu Asn Gln Glu Asp Gln Leu Val Trp Arg Thr Phe Pro Ser Val Lys  
65 70 75 80

Lys Phe Trp Ala Ser Pro Arg Gln Phe Ala Leu Gly His Trp Gly Lys  
                             85                            90                            95  
 Cys Arg Asp Asn Arg Gln Ala Lys Pro Leu Leu Ser Glu Glu Phe Phe  
                             100                            105                            110  
 5 Ala Thr Val Lys Glu Gly Tyr Gln Ile His Gln Asn Gln His Gln Gly  
                             115                            120                            125  
 Gln Ile Ile His Gly Asp Arg His Cys Arg Trp Gln Phe Thr Val Glu  
                             130                            135                            140  
 Pro Glu Val Thr Trp Gly Ser Pro Asn Arg Phe Pro Arg Ala Thr Ala  
 10 145                            150                            155                            160  
 Gly Trp Leu Ser Phe Leu Pro Leu Phe Asp Pro Gly Trp Gln Ile Leu  
                             165                            170                            175  
 Leu Ala Gln Gly Arg Ala His Gly Trp Leu Lys Trp Gln Arg Glu Gln  
                             180                            185                            190  
 15 Tyr Glu Phe Asp His Ala Leu Val Tyr Ala Glu Lys Asn Trp Gly His  
                             195                            200                            205  
 Ser Phe Pro Ser Arg Trp Phe Trp Leu Gln Ala Asn Tyr Phe Pro Asp  
                             210                            215                            220  
 His Pro Gly Leu Ser Val Thr Ala Ala Gly Gly Glu Arg Ile Val Leu  
 20 225                            230                            235                            240  
 Gly Arg Pro Glu Glu Val Ala Leu Ile Gly Leu His His Gln Gly Asn  
                             245                            250                            255  
 Phe Tyr Glu Phe Gly Pro Gly His Gly Thr Val Thr Trp Gln Val Ala  
                             260                            265                            270  
 25 Pro Trp Gly Arg Trp Gln Leu Lys Ala Ser Asn Asp Arg Tyr Trp Val  
                             275                            280                            285  
 Lys Leu Ser Gly Lys Thr Asp Lys Lys Gly Ser Leu Val His Thr Pro  
                             290                            295                            300  
 Thr Ala Gln Gly Leu Gln Leu Asn Cys Arg Asp Thr Thr Arg Gly Tyr  
 30 305                            310                            315                            320  
 Leu Tyr Leu Gln Leu Gly Ser Val Gly His Gly Leu Ile Val Gln Gly  
                             325                            330                            335  
 Glu Thr Asp Thr Ala Gly Leu Glu Val Gly Gly Asp Trp Gly Leu Thr  
                             340                            345                            350  
 35 Glu Glu Asn Leu Ser Lys Lys Thr Val Pro Phe  
                             355                            360

<210> 40  
 <211> 56  
 40 <212> DNA  
 <213> Artifical Sequence

<400> 40  
 cgcgatttaa atggcgcgcc ctgcaggcgg ccgcctgcag ggcgcgccat ttaaatt  
 45  
 <210> 41

56



<211> 32  
<212> DNA  
<213> Artificial Sequence

5 <400> 41  
tcgaggatcc gcggccgcaa gcttctctgca gg 32

<210> 42  
<211> 32  
10 <212> DNA  
<213> Artificial Sequence

<400> 42  
tcgacctgca ggaagcttgc ggccgcggat cc 32

15 <210> 43  
<211> 32  
<212> DNA  
<213> Artificial Sequence

20 <400> 43  
tcgacctgca ggaagcttgc ggccgcggat cc 32

<210> 44  
25 <211> 32  
<212> DNA  
<213> Artificial Sequence

<400> 44  
30 tcgaggatcc gcggccgcaa gcttctctgca gg 32

<210> 45  
<211> 36  
<212> DNA  
35 <213> Artificial Sequence

<400> 45  
tcgaggatcc gcggccgcaa gcttctctgca ggagct 36

40 <210> 46  
<211> 28  
<212> DNA  
<213> Artificial Sequence

45 <400> 46  
cctgcaggaa gcttgccggcc gcggatcc 28

<210> 47  
<211> 36  
<212> DNA  
5 <213> Artificial Sequence

<400> 47  
tcgacctgca ggaagcttgc ggccgcggat ccagct 36

10 <210> 48  
<211> 28  
<212> DNA  
<213> Artificial Sequence

15 <400> 48  
ggatccgcgg ccgcaagctt cctgcagg 28

<210> 49  
<211> 39  
20 <212> DNA  
<213> Artificial Sequence

<400> 49  
gatcacctgc aggaagcttg cggccgcgga tccaatgca 39

25 <210> 50  
<211> 31  
<212> DNA  
<213> Artificial Sequence

30 <400> 50  
ttggatccgc ggccgcaagc ttctgcagg t 31

<210> 51  
35 <211> 41  
<212> DNA  
<213> Artificial Sequence

<400> 51  
40 ggatccgcgg ccgcacaatg gagtctctgc tctctagttc t 41

<210> 52  
<211> 38  
<212> DNA  
45 <213> Artificial Sequence

<400> 52  
ggatcctgca ggtcacttca aaaaaggtaa cagcaagt 38

5 <210> 53  
<211> 45  
<212> DNA  
<213> Artifical Sequence

10 <400> 53  
ggatccgcgg ccgcacaatg gcgttttttg ggctctcccg tgttt 45

15 <210> 54  
<211> 40  
<212> DNA  
<213> Artifical Sequence

20 <400> 54  
ggatcctgca ggttattgaa aacttcttcc aagtacaact 40

25 <210> 55  
<211> 38  
<212> DNA  
<213> Artifical Sequence

30 <400> 55  
ggatccgcgg ccgcacaatg tggcgaagat ctgttggt 38

35 <210> 56  
<211> 37  
<212> DNA  
<213> Artifical Sequence

40 <400> 56  
ggatcctgca ggatcatggag agtagaagga aggagct 37

45 <210> 57  
<211> 50  
<212> DNA  
<213> Artifical Sequence

50 <400> 57  
ggatccgcgg ccgcacaatg gtacttgccg aggttccaaa gcttgccctt 50

55 <210> 58  
<211> 38  
<212> DNA

<213> Artificial Sequence

<400> 58  
5 ggatcctgca ggtcacttgt ttctgggtgat gactctat 38

<210> 59  
<211> 38  
<212> DNA  
<213> Artificial Sequence

10

<400> 59  
ggatccgcgg ccgcacaatg atttcgattc tcaacact 38

<210> 60  
15 <211> 36  
<212> DNA  
<213> Artificial Sequence

<400> 60  
20 ggatcctgca ggtcagtggt gcgatgctaa tgccgt 36

<210> 61  
<211> 22  
<212> DNA  
25 <213> Artificial Sequence

<400> 61  
taatgtgtac attgtcggcc tc 22

30 <210> 62  
<211> 60  
<212> DNA  
<213> Artificial Sequence

35 <400> 62  
gcaatgtaac atcagagatt ttgagacaca acgtggcttt ccacaattcc ccgcaccgtc 60

<210> 63  
<211> 22  
40 <212> DNA  
<213> Artificial Sequence

<400> 63  
aggctaataa gcacaaatgg ga 22

45 <210> 64

<211> 63  
 <212> DNA  
 <213> Artifical Sequence

5 <400> 64  
 ggtatgagtc agcaacacct tcttcacgag gcagacctca gcggaattgg tttagggttat 60  
 ccc 63

<210> 65  
 10 <211> 26  
 <212> DNA  
 <213> Artifical Sequence

<400> 65  
 15 ggatccatgg ttgcccacaaac cccatc 26

<210> 66  
 <211> 61  
 <212> DNA  
 20 <213> Artifical Sequence

<400> 66  
 gcaatgtaac atcagagatt ttgagacaca acgtggcttt gggtaagcaa caatgaccgg 60  
 c 61

25 <210> 67  
 <211> 25  
 <212> DNA  
 <213> Artifical Sequence

30 <400> 67  
 gaattctcaa agccagccca gtaac 25

<210> 68  
 35 <211> 63  
 <212> DNA  
 <213> Artifical Sequence

<400> 68  
 40 ggtatgagtc agcaacacct tcttcacgag gcagacctca gcgggtgcga aaagggtttt 60  
 ccc 63

<210> 69  
 <211> 23  
 45 <212> DNA  
 <213> Artifical Sequence

<400> 69  
ccagtgggttt aggctgtgtg gtc 23

5 <210> 70  
<211> 21  
<212> DNA  
<213> Artifical Sequence

10 <400> 70  
ctgagttgga tgtattggat c 21

<210> 71  
<211> 28

15 <212> DNA  
<213> Artifical Sequence

<400> 71  
ggatccatgg ttacttcgac aaaaatcc 28

20 <210> 72  
<211> 60  
<212> DNA  
<213> Artifical Sequence

25 <400> 72  
gcaatgtaac atcagagatt ttgagacaca acgtggcttt gctaggcaac cgcttagtac 60

<210> 73

30 <211> 28  
<212> DNA  
<213> Artifical Sequence

<400> 73

35 gaattcttaa cccaacagta aagttccc 28

<210> 74  
<211> 63  
<212> DNA

40 <213> Artifical Sequence

<400> 74  
ggatagagtc agcaacacct tcttcacgag gcagacctca gcgcggcat tgtcttttac 60  
atg 63

45 <210> 75

<211> 20  
<212> DNA  
<213> Artificial Sequence

5 <400> 75  
ggaacccttg cagccgcttc 20

<210> 76  
<211> 22  
10 <212> DNA  
<213> Artificial Sequence

<400> 76  
gtatgcccaa ctggtgcaga gg 22

15 <210> 77  
<211> 28  
<212> DNA  
<213> Artificial Sequence

20 <400> 77  
ggatccatgt ctgacacaca aaataccg 28

<210> 78  
25 <211> 62  
<212> DNA  
<213> Artificial Sequence

<400> 78  
30 gcaatgtaac atcagagatt ttgagacaca acgtggcttt cgccaatacc agccaccaac 60  
ag 62

<210> 79  
<211> 27  
35 <212> DNA  
<213> Artificial Sequence

<400> 79  
gaattctcaa atccccgcat ggcctag 27

40 <210> 80  
<211> 65  
<212> DNA  
<213> Artificial Sequence

45 <400> 80

ggtatgagtc agcaacacct tcttcacgag gcagacctca gcggcctacg gcttggacgt 60  
gtggg 65

5 <210> 81  
<211> 21  
<212> DNA  
<213> Artificial Sequence

10 <400> 81  
cacttggatt cccctgatct g 21

15 <210> 82  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<400> 82  
gcaatacccg cttggaaaac g 21

20 <210> 83  
<211> 29  
<212> DNA  
<213> Artificial Sequence

25 <400> 83  
ggatccatga ccgaatcttc gcccttagc 29

30 <210> 84  
<211> 61  
<212> DNA  
<213> Artificial Sequence

35 <400> 84  
gcaatgtaac atcagagatt ttgagacaca acgtggcttt caatcctagg tagccgaggc 60  
g 61

40 <210> 85  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<400> 85  
gaattcttag cccaggccag cccagcc 27

45 <210> 86  
<211> 66



<212> DNA  
<213> Artifical Sequence

<400> 86  
5 ggtatgagtc agcaacacct tcttcacgag gcagacctca gcggggaatt gatttggtta 60  
attacc 66

<210> 87  
<211> 21  
10 <212> DNA  
<213> Artifical Sequence

<400> 87  
15 gcgatcgcca ttatcgcttg g 21

<210> 88  
<211> 24  
<212> DNA  
<213> Artifical Sequence  
20

<400> 88  
gcagactggc aattatcagt aacg 24

<210> 89  
25 <211> 25  
<212> DNA  
<213> Artifical Sequence

<400> 89  
30 ccatggattc gagtaaagtt gtcgc 25

<210> 90  
<211> 0  
<213> Artifical Sequence  
35

<400> 90  
gaattcactt caaaaaaggt aacag

<210> 91  
40 <211> 4550  
<212> DNA  
<213> Arabidopsis sp

<400> 91  
45 attttacacc aatttgatca cttaactaaa ttaattaaat tagatgatta tcccaccata 60  
tttttgagca ttaaaccata aaaccatagt tataagtaac tgttttaatc gaatatgact 120

	cgattaagat	taggaaaaat	ttataaccgg	taattaagaa	aacattaacc	gtagtaaccg	180
	taaatgccga	ttcctccctt	gtctaaaaga	cagaaaacat	atattttatt	ttgccccata	240
	tgtttcactc	tatttaattt	caggcacaat	acttttggtt	ggtaacaaaa	ctaaaaagga	300
	caacacgtga	tacttttctt	cgtccgtcag	tcagattttt	tttaaactag	aaacaagtgg	360
5	caaatctaca	ccacattttt	tgcttaattt	attaacttgt	aagtttttaa	ttcctaaaaa	420
	agtctaacta	attcttctaa	tataagtaca	ttccctaaat	ttcccaaaaa	gtcaaattaa	480
	taattttcaa	aatctaattt	aaatatctaa	taattcaaaa	tcattaaaaa	gacacgcaac	540
	aatgacacca	attaatcatc	ctcgacccac	acaattctac	agttctcatg	ctaaaccata	600
	ttttttgtct	tctgttctct	caaaatcatt	tctttctctt	ctttgattcc	caaagatcac	660
10	ttctttgtct	ttgatttttg	attttttttt	tctctggcgt	gaaggaagaa	gcttttattt	720
	atggagtctc	tgctctctag	ttcttctctt	gtttccgctg	gtaaatctcg	tccttttctg	780
	gtttcaggtt	ttatttggtg	tttaggtttt	gtttttgtga	ttcagaacca	tacaaaaagt	840
	ttgaactttt	ctgaatataa	aataaggaaa	aagtttcgat	ttttataatg	aattgtttac	900
	tagatcgaag	taggtgacaa	aggttattgt	gtggagaagc	ataatttctg	ggcttgactt	960
15	tgaattttgt	ttctcatgca	tgcaacttat	caatcagctg	gtgggttttg	ttggaagaag	1020
	cagaatctaa	agctccactc	tttatcaggt	tcgttagggg	tttatggggt	tttgaaatta	1080
	aatactcaat	catcttagtc	tcattattct	attgggtgaa	tcacattttt	taatttgga	1140
	tttatgagac	aatgtatggt	ggacttaggt	gaagtctctt	tctttgggta	tagttgaagt	1200
	gttactgatg	ttgttttagct	ctttacacca	atatatacac	ccaattttgc	agaaatccga	1260
20	gttctgcgtt	gtgattcgag	taaagttgtc	gcaaaaccga	agtttaggaa	caatcttggt	1320
	aggcctgatg	gtcaaggatc	ttcattggtg	ttgtatccaa	aacataagtc	gagatttcgg	1380
	gttaatgcc	ctgcgggtca	gcctgaggct	ttcgactcga	atagcaaaca	gaagtctttt	1440
	agagactcgt	tagatgcgtt	ttacagggtt	tctaggcctc	atacagttat	tggcacaggt	1500
	aagtttctct	ttaaaaatgt	aactctttta	aaacgcaatc	tttcagggtt	ttcaaggaga	1560
25	taacattagc	tctgtgattg	gatttgcagg	tgcttagcat	tttatctgta	tctttcttag	1620
	cagtagagaa	ggtttctgat	atatctcctt	tacttttcac	tggcatcttg	gaggtaatga	1680
	atatataaca	cataatgacc	gatgaagaag	atacattttt	ttcgtctctc	tgtttaaaaca	1740
	attgggtttt	gttttcaggc	tggtgttgca	gctctcatga	tgaacattta	catagttggg	1800
	ctaaatcagc	tgtctgatgt	tgaaatagat	aaggtaacat	gcaaattttt	ttcatatgag	1860
30	ttcgagagac	tgatgagatt	aatagcagct	agtgcctaga	tcactctctat	gtgggttttt	1920
	gcaggttaac	aagccctatc	ttccattggc	atcaggagaa	tattctgtta	acaccggcat	1980
	tgcaatagta	gcttccctct	ccatcatggt	atggtgccat	tttcacaaaa	tttcaacttt	2040
	tagaattcta	taagttactg	aaatagtttg	ttataaatcg	ttatagagtt	tctggcttgg	2100
	gtggattggt	ggttcatggc	cattgttctg	ggctcttttt	gtgagtttca	tgctcggtac	2160
35	tgcatactct	atcaatgtaa	gtaagtttct	caatactaga	atttggttca	aatcaaaatc	2220
	tgcagtttct	agtttttaggt	taatgagggt	ttaataactt	acttctacta	caaacagttg	2280
	ccacttttac	ggtggaaaag	atttgcattg	gttgcagcaa	tgtgtatcct	cgctgtccga	2340
	gctattattg	ttcaaatcgc	cttttatcta	catattcagg	tactaaacca	ttttccttat	2400
	gttttgtagt	tgttttcatc	aaaatcactt	ttatattact	aaagctgtga	aactttgttg	2460
40	cagacacatg	tgtttggaag	accaatcttg	ttcactaggc	ctcttatttt	cgccactgcg	2520
	tttatgagct	ttttctctgt	cgttattgca	ttgtttaagg	taaacaaaga	tggaanaaga	2580
	ttaaatctat	gtatacttaa	agtaaagcat	tctactgtta	ttgatgagaa	gttttctttt	2640
	ttgggtggat	gcaggatata	cctgatatcg	aaggggataa	gatattcgga	atccgatcat	2700
	tctctgtaac	tctgggtcag	aaacgggtac	gatatctaaa	ctaaagaaat	tgttttgact	2760
45	caagtgttgg	attaagatta	cagaagaaag	aaaactgttt	ttgtttcttg	caaaattcag	2820
	gtgttttgga	catgtgttac	actacttcaa	atggcttacg	ctgttgcaat	tctagttgga	2880

	gccacatctc cattcatatg gagcaaagtc atctcggtaa caatctttct ttacccatcg	2940
	aaaactcgct aattcatcgt ttgagtggtg ctggtttcat tttgttccgt tctgttgatt	3000
	ttttttcagg ttgtgggtca tggtatactc gcaacaactt tgtgggctcg agctaagtcc	3060
	gttgatctga gtagcaaaac cgaaataact tcatgttata tgttcatatg gaagggtaga	3120
5	ttcgtttata aatagagttt ttactgcctt tttatgcgct ccaatttgga attaaaatag	3180
	cctttcagtt tcatcgaatc accattatac tgataaattc tcatttctgc atcagctctt	3240
	ttatgcagag tacttgctgt tacctttttt gaagtgactg acattagaag agaagaagat	3300
	ggagataaaa gaataagtca tcactatgct tctgttttta ttacaagttc atgaaattag	3360
	gtagtgaact agtgaattag agttttattc tgaaacatgg cagactgcaa aaatatgtca	3420
10	aagatatgaa tttctgttgg gtaaagaagt ctctgcttgg gcaaaatctt aagggttcggt	3480
	gtgttgatat aatgctaagc gaagaaatcg attctatgta gaaatttccg aaactatgtg	3540
	taaacatgtc agaacatctc cattctatat cttcttctgc aagaaagctc tgtttttatc	3600
	acctaaactc tttatctctg tgtagttaag atatgtatat gtacgtgact acattttttt	3660
	gttgatgtaa tttgcagaac gtatggattt ttgttagaaa gcatgagttc gaaagtatat	3720
15	gtttatatat atggataatt cagacctaac gtcgaagctc acaagcataa attcactact	3780
	atagtttgct ctgtaataga tagttccatt gatgtcttga aactgtacgt aactgcctgg	3840
	gcgttttgtg gttgatactg actactgagt gttctttgtg agtgttgtaa gtatacaaga	3900
	agaagaatat aggctcacgg gaacgactgt ggtggaagat gaaatggaga tcatcacgta	3960
	gcggctttgc caaagaccga gtcacgatcg agtctatgaa gtctttacag ctgctgatta	4020
20	tgattgacca ttgcttagag acgcattgga atcttactag ggacttgcct gggagtttct	4080
	tcaagtacgt gtcagatcat acgatgtagg agatttcacg gctttgatgt gtttgtttgg	4140
	agtcacaatg cttaatgggc ttattggccc aataatagct agctcttttg ctttagccgt	4200
	ttcgtttgtc ccttgggtgg gagtattatt agggtatggg gtgaccaaag tcaccagacc	4260
	tagagtgaat ctagtagagt cctagaccat ggtccatggc ttttatttgt aatttgaaaa	4320
25	atgaacaatt ctttttgtaa ggaaaacttt tatatagtag acgtttacta tatagaaact	4380
	agttgaacta acttcgtgca attgcataat aatgggtgtg aatagagggt gcaaaactca	4440
	ataaacattt cgacgtacca agagttcgaa acaataagca aatagattt ttttgcttca	4500
	gactaatttg tacaatgaat ggtaataaaa ccattgaagc ttttattaat	4550
30	<210> 92	
	<211> 4450	
	<212> DNA	
	<213> Arabidopsis sp	
35	<400> 92	
	tttaggttac aaaatcaatg atattgcgta tgtcaactat aaaagccaaa agtaaagcct	60
	cttgtttgac cagaagggtca tgatcattgt atacatacag ccaaactacc tcctggaaga	120
	aaagacatgg atcccaaaca acaacaatag cttcttttac aagaaccagt agtaactagt	180
	cactaatcta aaagagttta gtttcagctt ttctggcaat ggctccttga tcatttcaat	240
40	cctgaaggag acccactttg tagcaagacc atgtcctctg tttcacttac agtgtgtctc	300
	aaaagtctac ttcaattctt catatatagg ttcttcacac tacagcttca tcctcattcg	360
	ttgacagaga gagagtcttt attgaaaact tcttccaagt acaactccac taaatataat	420
	agcaccaaac cacttgttcg acacaaatct gtacagatat aaaaacacta ttaggttttc	480
	caaggcaaat cacataattg gattgtgaaa gagtacaaaa gataaaccce aattttcata	540
45	ctttctactg cagtcagcac cagatgataa gtcagctgtc cctatttgcc atcctaactg	600
	tcctgatgca gcggccagtg atgcgtaata ttgccaccct taatcattag agcgagaaac	660

	aaaaagaatc	aaaagacagt	aaatggaatt	aggaatcaca	aatgagtcct	tgtaaagttt	720
	attgagtacc	gagatctgca	ctgaatccag	aaagtgcag	aaaacctatg	gatgctgtgc	780
	caaatccagt	taaccaaagc	tttgtattat	caccgaatct	aagggctgtt	gacttaacac	840
	caacttttac	atcatcttct	ttgtcctgga	gacacaatat	attagacatt	agtccatgga	900
5	aaaaaaatga	tttaacctag	aatatctcaa	aattacttgc	ataaaaactg	aacttgagct	960
	gaaatttttg	gttcgtagct	tgtggcatat	actatttcat	tttcaatggg	ccacaaaggt	1020
	aactttcttt	tctcacttct	gttgcaaacg	ggaagacttt	tatggggcta	actcttcact	1080
	taaagtatag	aatcagatg	gaaaaggtgg	gagatcaggg	taattttctt	ctttatgatt	1140
	gacaaaagtc	gaacatcgaa	atggatgcat	ttgcatgaga	catgaaacaa	aagctgaaaa	1200
10	agaaatctgt	ggtgggtgaag	ctagaaaaag	aaaacaaagc	aagcaatatg	cacacattga	1260
	gattaactac	tttgctactg	gtcataatca	aatagatttt	gaagctaaaa	aataaaaaagt	1320
	gaatatacct	gatgtgcata	aatagtatca	taaacaaggg	tccagcagac	tccggagaga	1380
	tagagagggg	gtacaataga	tgggtgctatg	cttcctttta	ctgcagtgca	tcctaacaat	1440
	gctccccagt	ttatgggtcaa	acctaataag	gcttgaggct	gcaattataa	aaacgaatca	1500
15	atcataagaa	aatcagaaaa	tatataatgt	ctaactttga	gaagccagaa	tagattttaa	1560
	ttacccaaaa	tgtaaacctc	ttcataagtg	ggtaggaaaa	gacaagtaac	aaagatgaag	1620
	cccctaaaa	acggctgcag	aatatacata	ctgaaatgag	ctcaagtaga	aaagaatttg	1680
	atcacaaaa	taaagacaag	acctgagaac	atatcttcag	aatttgggca	aactacataa	1740
	gggtgaacca	tatgtgtatg	tgaattttta	aacaaacact	tgcaaatagc	cgacttttagg	1800
20	gcaagtaaaa	aatccaaaca	aacctgtaat	tgttaagttg	gagaagaatc	cctaagccta	1860
	aaagcaactg	cagcccgaga	aatccaatcc	cttgaaatgg	tgtcaaaaaga	ccactggcga	1920
	taggtcttag	ttttgtacga	tcaacctgga	tataaaagaa	atttgtaaga	caacataatc	1980
	taaaacaaaa	caaccataca	aaatcttgag	ctttacatac	aagcaaccca	tctttgttta	2040
	tggaagaatg	aatccagtta	catgaatgct	gtgtatctac	cctaactact	aaacacatat	2100
25	ttcaatcgaa	aaacatattc	caccttcacc	atatctaaca	cctgaagtct	ttcacttttt	2160
	gaacgaagtc	atcagaacat	gcagataagc	tattacccaa	aacagagata	tgactggaaa	2220
	tgttgctcgt	aattgatcca	acatagaaaa	atcaagacca	gttccagatg	tcaaagcaat	2280
	aacactttcc	caccatgggt	acagaaacca	tagttacaca	aaacatgttt	cctaaaccaa	2340
	cataactaaag	ggatatataa	atttgacatc	actttatcac	cataccataa	gatagcttaa	2400
30	aaacaaactg	acctttgtat	ctatgtcctg	atcaagcaga	tcatttatag	tacaaccagc	2460
	acctctaaga	agtaatgctc	cgcaacccaa	taaagccata	tattttaaac	ttggaaggct	2520
	tccaggatca	gcagccaacg	caatcgacct	atacaacaat	gatggagatt	cagagtatcg	2580
	atctatttac	atagctctgg	aactagatcc	atgacgaaac	atggaacatc	gttataatat	2640
	ctaaagactt	ccaaacagat	tcctgagtaa	gaaacccagt	ggaactatag	tactgtaaca	2700
35	tatataaaat	caaagaaaac	tcaggtttat	agcattatcc	aatcctgatt	tctgccaatc	2760
	cttaaccact	ctcccatgct	atcaaaaacc	tcagctcaag	atcatactac	ctaattgcct	2820
	atgagctctt	gggaagatca	ttatggattt	gataactgaa	aaaagtaaca	gagaaatagc	2880
	agactgcaag	aactactcca	aactttctcca	ctgatatgta	tgtagtctaa	caataataaa	2940
	cagacataaa	ttctttttatc	aagcttcaag	agcaagttag	tcagaaaaca	tcacagccaa	3000
40	accaaccagg	aaaacacata	actttatcac	ataaaaactaa	atttaaatgta	atctgactta	3060
	acataaacca	tcctttggga	cgaaaggaaa	ctatataaac	atgcagtctt	tctttccctc	3120
	agctattctt	tcggatggat	tataatgaat	ctcaaaagtg	aaatgtcttg	attctcagct	3180
	acattactca	aaggcgaaga	taaacttacc	acatacaagg	ccacgcaagc	aaccaagttc	3240
	caatggggtt	atccaatcga	gcaagcttag	cataacctct	aacttcttct	ggtaaataca	3300
45	aatctatcca	agaagcttcc	ttaacaacaa	caccatcact	cttctcctta	tcattctttct	3360
	tcggctttcc	ctccaaaacc	gaagaagacg	acgacattcc	acaaattaat	ctgtaattcc	3420

	aaccaacacc	aaaaaacttc	tcctgatgca	attctcttcc	tttactccat	acttggtaat	3480
	tatcattcca	tgaaggataa	cacttagtga	aaggatttgt	gtaatgggta	gtcacaggat	3540
	tggacaagga	tttatgttgt	gattgcaaaa	gagcagagga	agaagatgga	gttacggaga	3600
	cggaagattt	caacaaccgt	cttgaaacac	gggagagccc	aaaaaacgcc	atctttgaga	3660
5	gaaattgttg	cctggaagaa	acaaagactt	gagatttcaa	acgtaagtga	attcttacga	3720
	acgaaagcta	acttctcaag	agaatcagat	tagtgattcc	tcaaaaacaa	acaaaactat	3780
	ctaatttcag	tttcgagtga	tgaagcctta	agaatctaga	acctccatgg	cgtttcta	3840
	ctctcagaga	taatcgaatt	ccttaaacaa	tcaaagctta	gaaagagaag	aacaacaaca	3900
	acaacaaaaa	aatcagatt	aacaaccgac	cagagagcaa	cgacgacgcc	ggcgagaaag	3960
10	agcacgtcgt	ctcggagcaa	gacttcttct	ccagtaaccc	ggatggatcg	ttaatgggcc	4020
	tgtagattat	tatatattggg	ccgaaacaat	tgggtcagca	aaaacttggg	ggataatgaa	4080
	gaaacacgta	cagtatgcat	ttaggctcca	aattaattgg	ccatataatt	cgaatcagat	4140
	aaactaatca	accctacct	tacttatttc	tactgtttt	tatttctacc	ttagtagttg	4200
	aagaaacact	tttatattatc	ttttcgggac	ccaaatttga	taggatcggg	ccattactca	4260
15	tgagcgtcag	acacatatta	gccttatcag	attagtgggg	taagggtttt	ttaattcggg	4320
	aagaagcaac	aatcaatgtc	ggagaaatta	aagaatctgc	atgggcgtgg	cgtgatgata	4380
	tgtgcatatg	gagtcagttg	ccgatcatat	ataactattt	ataaactaca	tataaagact	4440
	actaatagat						4450
20	<210>	93					
	<211>	2850					
	<212>	DNA					
	<213>	Arabidopsis sp					
25	<400>	93					
	aattaaaatt	tgagcgggtct	aaaccattag	accgttttaga	gatccctcca	acccaaaata	60
	gtcgattttc	acgtcttgaa	catatatattg	gccttaattct	gtgtgggttag	taaagacttt	120
	tattgggtcaa	agaaaaacaa	ccatggccca	acatgttgat	acttttattt	aattatacaa	180
	gtacccttga	attctctgaa	atatatattga	ttgaccacga	tattaatttt	aattatcatt	240
30	tcctgtaaaa	gtgaaggagt	caccgtgact	cgtcgtaatc	tgaaaccaat	ctgttcatat	300
	gatgaagaag	tttctctcgt	tctcctccaa	cgcgtagaaa	attctgacgg	cttaacgatg	360
	tggcgaagat	ctgttggtta	tcgtttctct	tcaagaatct	ctgtttcttc	ttcgttacca	420
	aaccctagac	tgattccttg	gtcccgcgaa	ttatgtgccg	ttaatagctt	ctcccagcct	480
	ccggtctcga	cggaatcaac	tgctaagtta	gggatcactg	gtgttagatc	tgatgccaat	540
35	cgagtttttg	ccactgctac	tgccgccgct	acagctacag	ctaccaccgg	tgagatttcg	600
	tctagagttg	cggctttggc	tggattaggg	catcactacg	ctcgttgtta	ttgggagctt	660
	tctaaagcta	aacttaggta	tgtgtttact	tttcttttct	catgaaaaat	ctgaaaattt	720
	ccaattgttg	gattctttaa	ttctcatttg	ttttatgggt	gtagtatgct	tgtggttgca	780
	acttctggaa	ctgggtatat	tctgggtacg	ggaaatgctg	caattagctt	cccggggctt	840
40	tggttacacat	gtgcaggaac	catgatgatt	gctgcatctg	ctaattcctt	gaatcaggtc	900
	attgaaatgt	tgagaagttc	ataaattttcg	aatccttggt	gtgtttatgt	agttgatctt	960
	gcttgcttat	gtttatgtag	ttgaaaagtt	taaaaatttc	taatccttgg	tagttgatct	1020
	cgttggtttg	ttttttcatt	ttctagattt	ttgagataag	caatgattct	aagatgaaaa	1080
	gaacgatgct	aaggccattg	ccttcaggac	gtattagtgt	tccacacgct	gttgcatggg	1140
45	ctactattgc	tggtgcttct	ggtgcttggt	tgttggccag	caagggtgaat	gtttgttttt	1200
	ttatatgtga	tttctttgtt	ttatgaatgg	gtgattgaga	gattatggat	ctaaactttt	1260

	gcttccacga	caagggttatt	gcagactaat	atggttggtg	ctggacttgc	atctgccaat	1320
	cttgtacttt	atgcgtttgt	ttatactccg	ttgaagcaac	ttcacccctat	caatacatgg	1380
	gttggcgctg	ttggttggtg	tatcccaccc	ttgcttgggt	aaatttttgt	tccttttctt	1440
	ctttatttta	gcagattctg	ttttgttgga	tactgctttt	aattcaaaat	gtagtcatgg	1500
5	ttcaccaatt	ctatgcttat	ctattttgtg	tggtgtcagg	tgggcggcag	cgtctggtca	1560
	gatttcatac	aattcgatga	ttcttccagc	tgctctttac	ttttggcaga	tacctcattt	1620
	tatggccctt	gcacatctct	gccgcaatga	ttatgcagct	ggagggttaag	accatatggt	1680
	gtcatatgag	attagaatgt	ctccttccat	gtagtgttga	tcttgaacta	gttcaatttc	1740
	gtggaatgat	cagagtgtcc	tagatagtgt	cacagcagtc	gacatttttag	tggctagata	1800
10	atgagttctt	tccgttagag	ataaacattc	gcgaacattg	tttccagctt	ccgcgaccca	1860
	acttctgatt	ttgtttcttg	gtaccttggt	ttcagttaca	agatgttggt	actctttgat	1920
	ccgtcaggga	agagaatagc	agcagtggct	ctaaggaact	gctttttacat	gatccctctc	1980
	ggtttcatcg	cctatgactg	tgagtcttgt	agattcatct	tttttttgta	gtttattgac	2040
	tgcattgctg	tatctgattt	ttgctgttcc	ttccaatttt	tgtgacaggg	gggttaacct	2100
15	caagttgggt	ttgcctcgaa	tcaacacttc	tcacactagc	aatcgctgca	acagcatttt	2160
	cattctaccg	agaccggacc	atgcataaag	caaggaaaat	gttccatgcc	agtcttctct	2220
	tccttccctgt	tttcatgtct	ggtcttcttc	tacaccgtgt	ctctaattgat	aatcagcaac	2280
	aactcgtaga	agaagccgga	ttaacaaatt	ctgtatctgg	tgaagtcaaa	actcagaggc	2340
	gaaagaaacg	tgtggctcaa	cctccgggtg	cttatgcctc	tgctgcaccg	tttcccttcc	2400
20	tcccagctcc	ttccttctac	tctccatgat	aacctttaag	caagctattg	aatttttgga	2460
	aacagaaatt	aaaaaaaaaa	tctgaaaagt	tcttaagttt	aatctttggt	taataatgaa	2520
	gtggagaacg	catacaagtt	tatgtatttt	ttctcatctc	cacataattg	tattttttct	2580
	ctaagtatgt	ttcaaatgat	acaaaataca	tactttatca	attatctgat	caaattgatg	2640
	aatttttgag	ctttgacgtg	ttaggtctat	ctaataaacg	tagtaacgaa	tttggttttg	2700
25	gaaatgaaat	ccgataaccg	atgatggtgt	agagttaaac	gattaaaccg	ggttggttaa	2760
	aggtctcgag	tctcgacggc	tgcggaatc	ggaaaatcac	gattgaggac	tttgagctgc	2820
	cacgaagatg	gcgatgaggt	tgaaatcaat				2850

<210> 94  
 30 <211> 3660  
 <212> DNA  
 <213> Arabidopsis sp

<400> 94

35	tatttgtatt	tttattgtta	aatttttatga	tttcacccgg	tatatatcat	cccatattaa	60
	tattagattt	attttttggg	ctttattttgg	gttttctgatt	taaactgggc	ccattctgct	120
	tcaatgaaac	cctaattgggt	tttgtttggg	ctttggattt	aaaccgggccc	cattctgctt	180
	caatgaaggt	cctttgtcca	acaaaactaa	catccgacac	aactagtatt	gccaagagga	240
	togtgccaca	tggcagttat	tgaatcaaag	gccgcaaaa	ctgtaacgta	gacattactt	300
40	atctccggta	acggacaacc	actcgtttcc	cgaaacagca	actcacagac	tcacaccact	360
	ccagtctccg	gcttaactac	caccagagac	gattctctct	tccgtcgggt	ctatgacttc	420
	gattctcaac	actgtctcca	ccatccactc	ttccagagtt	acctccgtcg	atcgagtcgg	480
	agtcctctct	cttcggaatt	cggattccgt	tgagttcact	cgccggcggt	ctggtttctc	540
	gacgttgatc	tacgaatcac	ccggtagtta	gcattctggt	ggatagattg	atgaatgttt	600
45	tcttcgattt	tttttttact	gatcttggtg	tggatctctc	gtaggggcgga	gatttggtgt	660
	gcgtgcggcg	gagactgata	ctgataaagg	tatgattttt	tagttgtttt	tattttctct	720



	ctcttcaaaa	ttctcttttc	aaacactgtg	gcgtttgaat	ttccgacggc	agttaaactct	780
	cagacacctg	acaaggcacc	agccggtggt	tcaagcatta	accagcttct	cggtatcaaa	840
	ggagcatctc	aagaaactgt	aattttgttc	atctcctcag	aatcttttaa	attatcatat	900
	ttgtggataa	tgatgtgtta	gttttaggaat	tttcctacta	aaggtaatct	cttttgagga	960
5	caagtcttgt	tttttagctta	gaaatgatgt	gaaaatgttg	tttgtttagct	aaaaagagtt	1020
	tggtgttata	ttctgtattc	agaataaatg	gaagattcgt	cttcagctta	caaaaccagt	1080
	cacttggcct	ccactgggtt	ggggagtcgt	ctgtgggtgct	gctgcttcag	gtaatcatac	1140
	gaacctcttt	tggatcatgc	aatactgtac	agaaagtttt	ttcattttcc	ttccaattgt	1200
	ttcttctggc	agggaaacttt	cattggaccc	cagaggatgt	tgctaagtcg	attctttgca	1260
10	tgatgatgtc	tggtccttgt	cttactggct	atacacaggt	ctggttttac	acaacaaaaa	1320
	gctgacttgt	tcttattcta	gtgcatttgc	ttgggtgctac	aataacctag	acttgtcgat	1380
	ttccagacaa	tcaacgactg	gtatgataga	gatatcgacg	caattaatga	gccatatcgt	1440
	ccaattccat	ctggagcaat	atcagagcca	gaggttaactg	agacagaaca	ttgtgagctt	1500
	ttatctcttt	tgtgattctg	atttctcctt	actccttaaa	atgcagggtta	ttacacaagt	1560
15	ctgggtgcta	ttattgggag	gtccttggtat	tgctggaata	ttagatgtgt	gggtaagttg	1620
	gcccttctga	cattaactag	tacagttaaa	gggcacatca	gatttgctaa	aatcttcctt	1680
	tatcaggcag	ggcataccac	tcccactgtc	ttctatcttg	ctttgggagg	atcattgcta	1740
	tcttatatat	actctgctcc	acctcttaag	gtaagtttta	ttcctaactt	ccactctcta	1800
	gtgataagac	actccatcca	agttttggag	ttttgaatat	cgatatctga	actgatctca	1860
20	ttgcagctaa	aacaaaatgg	atgggttgga	aattttgcac	ttggagcaag	ctatattagt	1920
	ttgccatggt	aagatatctc	gtgtatcaat	aatatatggc	gttgttctca	tctcattgat	1980
	ttgtttcttg	ctcacttgac	tgatagggtg	gctggccaag	cattgtttgg	cactcttacg	2040
	ccagatgttg	ttgttctaac	actcttgtag	agcatagctg	gggtactctt	ttggcaaacc	2100
	ttttatgttg	cttttttctg	tatctgttgt	aatatgctct	tgcttcatgt	tgtacctttg	2160
25	tgataatgca	gttaggaata	gccattgtta	acgacttcaa	aagtgttgaa	ggagatagag	2220
	cattaggact	tcagtctctc	ccagtagctt	ttggcaccca	aactgcaaaa	tggatatgag	2280
	ttgggtgctat	agacattact	cagctttctg	ttgccggtat	gtactatcca	ctgtttttgt	2340
	gcagctgtgg	cttctatttc	ttttccttga	tcttatcaac	tggatattca	ccaatggtaa	2400
	agcacaat	aatgaagctg	aatcaacaaa	ggcaaaacat	aaaagtacat	tctaataaaa	2460
30	tgagctaata	aagaggaggc	atctactttt	atgtttcatt	agtgtgattg	atggattttc	2520
	atttcatgct	tctaaaacaa	gtattttcaa	cagtgtcatg	aaataacaga	acttatatct	2580
	tcatttgtac	ttttactagt	ggatgagtta	cacaatcatt	gttatagaac	caaatcaaag	2640
	gtagagatca	tcattagtat	atgtctatatt	tggttgagag	atatctatta	gcattctggga	2700
	aaccttatta	tgcgttggcg	ttgggttgctt	tgatcattcc	tcagattgtg	ttccaggtaa	2760
35	agacgttaac	agtctcacat	tataattaat	caaattcttg	tcactcgtct	gatttgctaca	2820
	ctcgcttcta	taaactgcag	tttaaatact	ttctcaagga	ccctgtcaaa	tacgacgtca	2880
	agtaccagg	aagtcaactt	agtacacatg	tttgtgttct	tttgaaatat	ctttgagagg	2940
	tctcttaate	agaagttgct	tgaacactc	atcttgatta	caggcaagcg	cgcagccatt	3000
	cttgggtgctc	ggaatatattg	taacggcatt	agcatcgcaa	cactgaaaaa	ggcgtatttt	3060
40	gatgggggtt	tgtcgaaagc	agagggtgtg	acacatcaaa	tgtggggcaag	tgatggcatc	3120
	aactagttta	aaagattttg	taaaatgtat	gtaccgttat	tactagaaac	aactcctgtt	3180
	gtatcaattt	agcaaaacgg	ctgagaaatt	gtaattgatg	ttaccgtatt	tgcgctccat	3240
	ttttgcattt	cctgctcata	tcgaggattg	gggtttatgt	tagttctgtc	acttctctgc	3300
	tttcagaatg	tttttgtttt	ctgtagtgga	ttttaactat	tttcatcact	ttttgtattg	3360
45	attctaaaca	tgtatccaca	taaaaacagt	aataataaaa	aatgataact	cctcaaaact	3420
	tttataatct	aaatctaaca	actagctagt	aaccaacta	acttcataca	attaatttga	3480

gaaactacaa	agactagact	atacatatgt	tatttaacaa	cttgaaactg	tggtattact	3540
acctgatttt	tttctattct	acagccattt	gatatgctgc	aatcttaaca	tatcaagtct	3600
cacgttggtg	gacacaacat	actatcacaa	gtaagacacg	aagtaaaacc	aaccggcaac	3660



(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
26 October 2000 (26.10.2000)

PCT

(10) International Publication Number  
**WO 00/63391 A3**

(51) International Patent Classification<sup>7</sup>: C12N 15/54,  
15/82, 9/10, 5/00, C12P 17/06

POST-BEITTENMILLER, Dusty [US/US]; 800 N.  
Lindbergh Blvd., St. Louis, MO 63167 (US).

(21) International Application Number: PCT/US00/10368

(74) Agent: RAE-VENTER LAW GROUP, P.C.; P.O. Box  
60039, Palo Alto, CA 94306 (US).

(22) International Filing Date: 14 April 2000 (14.04.2000)

(25) Filing Language: English

(26) Publication Language: English

(81) Designated States (*national*): AL, AM, AT, AU, AZ, BA,  
BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES,  
FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG,  
MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE,  
SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN,  
YU, ZW.

(30) Priority Data:  
60/129,899 15 April 1999 (15.04.1999) US  
60/146,461 30 July 1999 (30.07.1999) US

(84) Designated States (*regional*): European patent (AT, BE,  
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,  
NL, PT, SE).

(71) Applicant (*for all designated States except US*): CAL-  
GENE LLC [US/US]; 1920 Fifth Street, Davis, CA 95616  
(US).

Published:  
— with international search report

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): SAVIDGE, Beth  
[US/US]; 1920 Fifth Street, Davis, CA 95616 (US).  
LASSNER, Michael, W. [US/US]; 1920 Fifth Street,  
Davis, CA 95616 (US). WEISS, James, D. [US/US];  
800 N. Lindbergh Blvd., St. Louis, MO 63167 (US).

(88) Date of publication of the international search report:  
17 January 2002

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

(54) Title: NUCLEIC ACID SEQUENCES TO PROTEINS INVOLVED IN TOCOPHEROL SYNTHESIS

(57) Abstract: Nucleic acid sequences and methods are provided for producing plants and seeds having altered tocopherol content and compositions. The methods find particular use in increasing the tocopherol levels in plants, and in providing desirable tocopherol compositions in a host plant cell.

# INTERNATIONAL SEARCH REPORT

Inter. Application No

PCT/US 00/10368

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/54 C12N15/82 C12N9/10 C12N5/00 C12P17/06

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BIOSIS, WPI Data, EPO-Internal, EMBL

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL [Online] ACCESSION NO: AC003673, 11 December 1997 (1997-12-11) LIN, X., ET AL.: "Arabidopsis thaliana chromosome II section 110 of 255 of the complete sequence. Sequence from clones MSF3, F19F24." XP002153685 nts40740-43320	1-6, 13-16,18
X	--- DATABASE EMBL [Online] ACCESSION NO: AL035394, 9 February 1999 (1999-02-09) BEVAN, M., ET AL.: "Arabidopsis thaliana DNA chromosome 4, BAC clone F9D16 (ESSAII project)" XP002153686 nts 46219-49152 --- -/--	1-6, 13-16,18

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

12 June 2001

Date of mailing of the international search report

15.06.01

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Maddox, A

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 00/10368

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL [Online]  ACCESSION NO: B24116,  13 October 1997 (1997-10-13)  ROUNSLEY, S.D., ET AL.: "F18L14TF IGF  Arabidopsis thaliana genomic clone F18L14,  genomic survey sequence."  XP002153687  the whole document</p>	1-6
X	<p>---  DATABASE EMBL [Online]  ACCESSION NO: AC003672,  11 December 1997 (1997-12-11)  LIN, X., ET AL.: "Arabidopsis thaliana  chromosome II section 239 of 255 of the  complete sequence. Sequence from clones  F4I1, F16B22."  XP002153688  nts 363-2540</p>	1-6, 13-16,18
X	<p>---  DATABASE EMBL [Online]  ACCESSION NO: B29398,  13 October 1997 (1997-10-13)  ROUNSLEY, S.D., ET AL.: "F16B22TRC IGF  Arabidopsis thaliana genomic clone F16B22,  genomic survey sequence."  XP002153689  the whole document</p>	1-6
X	<p>---  DATABASE EMBL [Online]  ACCESSION NO: R30625,  11 August 1995 (1995-08-11)  NEWMAN, T., ET AL.: "13230 Lambda-PRL2  Arabidopsis thaliana cDNA clone 166L10T7,  mRNA sequence."  XP002153690  abstract</p>	1-6
X	<p>---  GAUBIER PASCALE ET AL: "A chlorophyll  synthetase gene from Arabidopsis  thaliana."  MOLECULAR &amp; GENERAL GENETICS,  vol. 249, no. 1, 1995, pages 58-64,  XP002153682  ISSN: 0026-8925  the whole document  -&amp; DATABASE TREMBL [Online]  ACCESSION NO: Q38833,  1 November 1996 (1996-11-01)  GAUBIER, P., ET AL.: "PUTATIVE CHLOROPHYLL  SYNTHETASE (G4)."  XP002169117  the whole document</p> <p>---</p> <p>-/--</p>	1-6, 13-16,18

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 00/10368

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE BIOSIS [Online]  BIOSCIENCES INFORMATION SERVICE,  PHILADELPHIA, PA, US;  October 1997 (1997-10)  OSTER U ET AL: "The G4 gene of Arabidopsis thaliana encodes a chlorophyll synthase of etiolated plants."  Database accession no. PREV199800047824  XP002153691  abstract  &amp; BOTANICA ACTA,  vol. 110, no. 5, October 1997 (1997-10),  pages 420-423,  ISSN: 0932-8629</p> <p>---</p>	1-5, 13-16
X	<p>LOPEZ J ET AL: "Sequence of the bch6 gene from Chloroflexus aurantiacus: Relationship between Chlorophyll synthase and other polyprenyltransferases"  JOURNAL OF BACTERIOLOGY, WASHINGTON, DC, US,  vol. 178, no. 11, 1996, pages 3369-3373,  XP002146399  ISSN: 0021-9193  the whole document</p> <p>---</p>	1-4
X	<p>DATABASE EMBL [Online]  ACCESSION NO: AC004077,  3 February 1998 (1998-02-03)  LIN, X., ET AL.: "Arabidopsis thaliana chromosome II section 190 of 255 of the complete sequence. Sequence from clones F13P17, T31E10."  XP002169118  /gene="At2g34630"  -&amp; DATABASE TREMBL [Online]  ACCESSION NO: 064684,  1 August 1998 (1998-08-01)  ROUNSLEY S.D., ET AL.: "T31E10.3 PROTEIN"  XP002169119  abstract</p> <p>---</p>	1-6, 13-16,18
X	<p>DATABASE EMBL [Online]  ACCESSION NO: T44803,  4 February 1995 (1995-02-04)  NEWMAN, T. ET AL.: "8066 Lambda-PRL2 Arabidopsis thaliana cDNA clone 124L9T7, mRNA sequence."  XP002169120  the whole document</p> <p>---</p>	1-5

-/--

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 00/10368

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL [Online]  ACCESSION NO: Z34566,  25 June 1994 (1994-06-25)  DESPREZ, T., ET AL.: "A. thaliana  transcribed sequence; clone VBVC03; 5'  end; Similar to Cytochrome c554 ;  Chloroflexus aurantiacus."  XP002169121  the whole document</p>	1-5
X	<p style="text-align: center;">---</p> <p>ZHU XUFEN ET AL: "Geranylgeranyl  pyrophosphate synthase encoded by the  newly isolated gene GGPS6 from Arabidopsis  thaliana is localized in mitochondria."  PLANT MOLECULAR BIOLOGY,  vol. 35, no. 3, 1997, pages 331-341,  XP002153683  ISSN: 0167-4412  the whole document</p>	1-5, 13-16, 29,30
X	<p style="text-align: center;">---</p> <p>DATABASE EMBL [Online]  ACCESSION NO: L40577,  15 April 1995 (1995-04-15)  SCOLNIK, P.A., ET AL.: "Arabidopsis  thaliana geranylgeranyl pyrophosphate  synthase-related protein mRNA, complete  cds."  XP002153692  the whole document</p>	1-5
X	<p style="text-align: center;">---</p> <p>CHUN P L ET AL: "Identification of a  maize endosperm-specific cDNA encoding  farnesyl pyrophosphate synthetase"  GENE,NL,ELSEVIER BIOMEDICAL PRESS.  AMSTERDAM,  vol. 171, no. 2, 1 June 1996 (1996-06-01),  pages 193-196; XP004042793  ISSN: 0378-1119  the whole document</p>	1-4,7
E	<p style="text-align: center;">---</p> <p>EP 1 033 405 A (CERES INC)  6 September 2000 (2000-09-06)  see SEQ ID NOS:34834-34836,38169  38171,50712,50713</p>	1-7, 13-16,18
E	<p style="text-align: center;">---</p> <p>WO 00 68393 A (PIONEER HI-BRED)  16 November 2000 (2000-11-16)</p> <p>see SEQ ID NOS: 1,2,3,4,9-14,21,22,23</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	1-16, 18-20, 22-25, 27-30, 32,33

# INTERNATIONAL SEARCH REPORT

Interr. Application No  
PCT/US 00/10368

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 00 14207 A (DU PONT ;MIAO GUO HUA (US); POWELL WAYNE (US); CAHOON REBECCA E (U) 16 March 2000 (2000-03-16) SEQ ID NOS:1,2,5,6,7,8,11 and 12 ---	1-4,7,9
P,X	DATABASE EMBL [Online] ACCESSION NO: AI795655, 7 July 1999 (1999-07-07) WALBOT, V.: "614004H08.x4 614 - root cDNA library from Walbot Lab Zea mays cDNA, mRNA sequence." XP002169131 the whole document ---	1-4,7
P,X	DATABASE EMBL [Online] ACCESSION NO: AI988542, 7 September 1999 (1999-09-07) SHOEMAKER, R., ET AL.: "sd03g09.y1 Gm-c1020 Glycine max cDNA clone GENOME SYSTEMS CLONE ID:Gm-c1020-665 5' similar to TR:064886 064886 PUTATIVE HEME A:FARNESYLTRANSFERASE. ;, mRNA sequence." XP002169132 the whole document ---	1-4,9
P,X	DATABASE EMBL [Online] ACCESSION NO: AI938569, 3 August 1999 (1999-08-03) SHOEMAKER, R., ET AL.: "sb55e11.y1 Gm-c1018 Glycine max cDNA clone GENOME SYSTEMS CLONE ID:Gm-c1018-69 5' similar to TR:064625 064625 F19F24.15 PROTEIN. ;, mRNA sequence." XP002169133 the whole document ---	1-4,9
P,X	DATABASE EMBL [Online] ACCESSION NO: AW306617, 21 January 2000 (2000-01-21) SHOEMAKER R., ET AL.: "se53b09.y1 Gm-c1017 Glycine max cDNA clone GENOME SYSTEMS CLONE ID:Gm-c1017-2610 5' similar to TR:064625 064625 F19F24.15 PROTEIN. ;, mRNA sequence" XP002169134 the whole document ---	1-4,9

-/--

# INTERNATIONAL SEARCH REPORT

Interr. Application No  
PCT/US 00/10368

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>DATABASE EMBL [Online]  ACCESSION NO: AI748688,  29 June 1999 (1999-06-29)  SHOEMAKER R.: "sb60f03.y1 Gm-cl010 Glycine  max cDNA clone GENOME SYSTEMS CLONE  ID:Gm-cl010-150 5' similar to TR:P73726  P73726 HYPOTHETICAL 34.4 KD PROTEIN.;,  mRNA sequence."  XP002169135  the whole document</p>	1-4,9
X	<p>---  DATABASE EMBL [Online]  ACCESSION NO: D64006,  30 September 1995 (1995-09-30)  TABATA, S., ET AL.: "Synechocystis sp.  PCC6803 complete genome, 25/27,  3138604-3270709."  XP002169122  nts 90109-90987  -&amp; DATABASE TREMBL [Online]  ACCESSION NO: Q55500,  1 November 1996 (1996-11-01)  TABATA, S., :  "4-HYDROXYBENZOATE-OCTAPRENYL  TRANSFERASE."  XP002169123  the whole document</p>	1-4,11, 12
X	<p>---  DATABASE EMBL [Online]  ACCESSION NO: D13960,  28 March 1996 (1996-03-28)  MURATA N.; ET AL.: "Synechocystis sp.  genes for heme O synthase and  virginiamycin acetyltransferase, complete  cds."  XP002169124  the whole document  -&amp; DATABASE TREMBL [Online]  ACCESSION NO: Q55207,  1 November 1996 (1996-11-01)  "CYTOSHROME C OXIDASE FOLDING PROTEIN"  XP002169125  abstract</p> <p style="text-align: center;">--- -/--</p>	1-4,11, 12

# INTERNATIONAL SEARCH REPORT

Interr. Application No  
PCT/US 00/10368

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL [Online]  ACCESSION NO: D64001,  30 September 1995 (1995-09-30)  TABATA, S., ET AL.: "Synechocystis sp.  PCC6803 complete genome, 20/27,  2539000-2644794."  XP002169126  nts 39188..40162  -&amp; DATABASE TREMBL [Online]  ACCESSION NO: Q55145,  1 November 1996 (1996-11-01)  TABATA, S.: "CHLOROPHYLL SYNTHASE 33 KDA  SUBUNIT."  XP002169127  the whole document</p>	1-4,11, 12
X	<p>---  DATABASE EMBL [Online]  ACCESSION NO: D90911,  31 October 1996 (1996-10-31)  TABATA S.;: "Synechocystis sp. PCC6803  complete genome, 13/27, 1576593-1719643."  XP002169128  nts 33234..34157  -&amp; DATABASE TREMBL [Online]  ACCESSION NO: P73962,  15 July 1998 (1998-07-15)  KANEKO, T., ET AL.: "PROBABLE  1,4-DIHYDROXY-2-NAPHTHOATE  OCTAPRENYLTRANSFERASE (EC 2.5.1.-)"  XP002169129  abstract</p>	1-4,11, 12
X	<p>---  DATABASE EMBL [Online]  ACCESSION NO: D90909,  31 October 1996 (1996-10-31)  TABATA, S.: "Synechocystis sp. PCC6803  complete genome, 11/27, 1311235-1430418."  XP002169130  nts 11453..12379 and 12438..13529  -&amp; DATABASE TREMBL [Online]  ACCESSION NO: P73726,  1 February 1997 (1997-02-01)  KANEKO, T., ET AL.: "HYPOTHETICAL 34.4 KDA  PROTEIN."  XP002169263  abstract  -&amp; DATABASE TREMBL [Online]  ACCESSION NO: P73727,  1 February 1997 (1997-02-01).  KANEKO, T., ET AL.: "HYPOTHETICAL 41.5 KDA  PROTEIN."  XP002169264  abstract</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	1-4,11, 12



# INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/US 00/10368

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 99 07867 A (CALGENE LLC) 18 February 1999 (1999-02-18)</p> <p>page 11, line 19 - line 26 ---</p>	<p>1-4, 13-15, 18-20, 22-25, 27-30, 32,33</p>
X	<p>WO 98 06862 A (SHEWMAKER CHRISTINE K ;CALGENE INC (US)) 19 February 1998 (1998-02-19)</p>	<p>1-4, 13-15, 18,29, 30,32,33</p>
A	<p>the whole document</p>	<p>19,20, 22-25, 27,28</p>
E	<p>---</p> <p>WO 00 61771 A (MONSANTO CO) 19 October 2000 (2000-10-19)</p> <p>page 81 -page 83 page 106 -page 107</p>	<p>1-4, 13-15, 18-20, 22-25, 27-30, 32,33</p>
A	<p>---</p> <p>NORRIS S R ET AL: "GENETIC DISSECTION OF CAROTENOID SYNTHESIS IN ARABIDOPSIS DEFINES PLASTOQUINONE AS AN ESSENTIAL COMPONENT OF PHYTOENE DESATURATION" PLANT CELL,US,AMERICAN SOCIETY OF PLANT PHYSIOLOGISTS, ROCKVILLE, MD, vol. 7, 1 December 1995 (1995-12-01), pages 2139-2149, XP002041909 ISSN: 1040-4651 the whole document</p>	<p>1-6, 13-16, 18-20, 22-25, 27-30, 32,33</p>
X	<p>---</p> <p>KUNTZ, M., ET AL.: "Identification of a cDNA for the plastid-located geranylgeranyl pyrophosphate synthase from Capsicum annuum: correlative increase in enzyme activity and transcript level during fruit ripening" THE PLANT JOURNAL, vol. 2, no. 1, 1992, XP002153684 the whole document</p>	<p>1-4, 13-15, 18,29,30</p>
X	<p>---</p> <p>US 5 876 964 A (GERSHENZON JONATHAN ET AL) 2 March 1999 (1999-03-02)</p> <p>the whole document</p>	<p>1-4, 13-15, 18,29,30</p>
X	<p>---</p> <p>US 5 545 816 A (AUSICH RODNEY L ET AL) 13 August 1996 (1996-08-13)</p> <p>the whole document</p> <p>---</p>	<p>1-4, 13-15, 18,29,30</p>
	-/--	

# INTERNATIONAL SEARCH REPORT

Interr. Application No  
PCT/US 00/10368

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 763 542 A (TOYOTA MOTOR CO LTD) 19 March 1997 (1997-03-19) the whole document ---	29,30
X	EP 0 674 000 A (TOYOTA MOTOR CO LTD) 27 September 1995 (1995-09-27) the whole document ---	29,30
P,X	WO 00 01650 A (DCV INC) 13 January 2000 (2000-01-13) the whole document ---	1-4, 13-15, 18,29,30
E	EP 1 063 297 A (KOREA KUMHO PETROCHEM CO LTD) 27 December 2000 (2000-12-27) the whole document ---	1-4, 13-15, 18,29,30
A	WO 97 27285 A (UNIV ARIZONA) 31 July 1997 (1997-07-31) the whole document ---	19,20, 22-25, 27-30, 32,33
A	WO 99 04622 A (UNIV NEVADA) 4 February 1999 (1999-02-04) the whole document ---	19,20, 22-25, 27-30, 32,33
X	WO 99 06580 A (BONETTA DARIO ;MCCOURT PETER (CA); GHASSEMIAN MAJID (CA); PERFORMA) 11 February 1999 (1999-02-11) claims 18,19 ---	1-5, 13-16,18
E	WO 00 22150 A (PIONEER HI BRED INT ;YALPANI NASSER (US); MEYER TERRY EUCLAIRE (US) 20 April 2000 (2000-04-20) the whole document ---	1-4, 13-15,18
E	WO 01 21650 A (COLDREN CHRIS ;DU PONT (US); WANG HONG (US); FLINT DENNIS (US); HA) 29 March 2001 (2001-03-29) the whole document -----	1-4, 13-16,18

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 00/10368

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
**Claims 8 and 10 are inconsistent with figs 2,3,and 9, since said figures do not make reference to sequence data. Said claims have been assumed as relating to SEQ ID NOS:19-31 and the prior art search has been made accordingly.**
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-6,13-16, and 18 all partially

Nucleic acid sequences encoding Arabidopsis aromatic prenyltransferases as defined by SEQ ID NOS:1-6 and constructs based on said sequences.

2. Claims: 1-6,13-16, and 18 all partially.

Nucleic acid sequences encoding Arabidopsis straight chain prenyltransferases as defined by SEQ ID NOS:11,12,16,17 and constructs based on said sequences.

3. Claims: 1-4,13-16 and 18 all partially,  
and 7 and 8 both completely

Nucleic acid sequences encoding corn prenyltransferases and constructs based on said sequences.

4. Claims: 1-4,13-16, and 18 all partially,  
and 9 and 10 both completely

Nucleic acid sequences encoding soybean prenyltransferases and constructs based on said sequences.

5. Claims: 1-4,13,14, and 18 all partially, and 11,12,  
and 17 all completely

Nucleic acid sequences encoding synechocystis prenyltransferases and constructs based on said sequences.

6. Claims: 19-33 all completely

Methods for production of tocopherols, increasing flux to tocopherol production in a host cell by transforming with a prenyltransferase nucleic acid sequence

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/10368

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 1033405 A	06-09-2000	NONE	
-----	-----	-----	-----
WO 0068393 A	16-11-2000	AU 4498500 A	21-11-2000
-----	-----	-----	-----
WO 0014207 A	16-03-2000	AU 5812199 A	27-03-2000
-----	-----	-----	-----
WO 9907867 A	18-02-1999	AU 8900298 A	01-03-1999
		CN 1275166 T	29-11-2000
		EP 1002117 A	24-05-2000
-----	-----	-----	-----
WO 9806862 A	19-02-1998	AU 4058497 A	06-03-1998
		BR 9713462 A	28-03-2000
		CN 1227609 A	01-09-1999
		EP 0925366 A	30-06-1999
-----	-----	-----	-----
WO 0061771 A	19-10-2000	AU 4231600 A	14-11-2000
-----	-----	-----	-----
US 5876964 A	02-03-1999	AU 1089099 A	03-05-1999
		EP 1023436 A	02-08-2000
		WO 0129188 A	26-04-2001
		WO 9919460 A	22-04-1999
-----	-----	-----	-----
US 5545816 A	13-08-1996	US 5618988 A	08-04-1997
		CA 2055447 A	03-09-1991
		EP 0471056 A	19-02-1992
		JP 5504686 T	22-07-1993
		WO 9113078 A	05-09-1991
		US 5530188 A	25-06-1996
		US 5530189 A	25-06-1996
		US 5684238 A	04-11-1997
		US 5656472 A	12-08-1997
-----	-----	-----	-----
EP 0763542 A	19-03-1997	JP 9065878 A	11-03-1997
		DE 69604994 D	09-12-1999
		DE 69604994 T	27-04-2000
		US 5882909 A	16-03-1999
		US 5885810 A	23-03-1999
		US 5807725 A	15-09-1998
-----	-----	-----	-----
EP 0674000 A	27-09-1995	JP 7308193 A	28-11-1995
		US 5773273 A	30-06-1998
-----	-----	-----	-----
WO 0001650 A	13-01-2000	AU 4863099 A	24-01-2000
		EP 1095002 A	02-05-2001
-----	-----	-----	-----
EP 1063297 A	27-12-2000	JP 2001000192 A	09-01-2001
-----	-----	-----	-----
WO 9727285 A	31-07-1997	US 6087563 A	11-07-2000
		AU 1845397 A	20-08-1997
		BR 9707200 A	28-12-1999
		EP 0877793 A	18-11-1998
		JP 11510708 T	21-09-1999
-----	-----	-----	-----
WO 9904622 A	04-02-1999	AU 8506198 A	16-02-1999
		EP 1009812 A	21-06-2000
-----	-----	-----	-----
WO 9906580 A	11-02-1999	AU 8598998 A	22-02-1999
		EP 1002116 A	24-05-2000

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Interr. Application No

PCT/US 00/10368

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9906580 A		ZA 9806872 A	02-02-1999
-----		-----	-----
WO 0022150 A	20-04-2000	AU 6290099 A	01-05-2000
-----		-----	-----
WO 0121650 A	29-03-2001	NONE	
-----		-----	-----